

Please type a plus sign (+) inside this box ☒

**UTILITY
PATENT APPLICATION
TRANSMITTAL**

Attorney Docket No.

ORT-1238

First Named Inventor or Application Identifier

Han-Cheng Zhang, Bruce E. Maryanoff, Anjali Pandey and Robert M. Scarborough

Express Mail Label No.

EL327264653US

(only for new nonprovisional applications under 37 CFR 1.53(b))

APPLICATION ELEMENTS

ADDRESS TO:

Assistant Commissioner for Patents
Box Patent Application
Washington, DC 20231

See MPEP Chapter 600 concerning utility patent application contents.

1. ☒ Fee Transmittal Form (attached hereto in duplicate)

2. ☒ Specification [Total Pages 51]

(Preferred arrangement set forth below)

- Descriptive Title of the Invention
- Cross References to Related Applications
- Statement Regarding Fed sponsored R&D
- Reference to Microfiche Appendix
- Background of the Invention
- Brief Summary of the Invention
- Brief Description of the Drawings (if filed)
- Detailed Description
- Claim(s)
- Abstract of the Disclosure

3. ☐ Drawing(s) (35 USC 113) [Total Sheets]

4. Oath or Declaration

a. ☒ Newly executed (original or copy)

b. ☐ Unexecuted original

c. ☐ Copy from a prior application (37 CFR 1.63(d))
(for continuation/divisional check boxes 5 and 16)

i. ☐ Deletion of Inventor(s)

Signed statement attached deleting inventor(s) named in the prior application, see 37 CFR 1.63(d)(2) and 1.33(b).

5. ☐ Incorporation by Reference
(useable if Box 4c is checked)

The entire disclosure of the prior application, from which a copy of the oath or declaration is supplied under Box 4c, is considered as being part of the disclosure of the accompanying application and is hereby incorporated by reference therein.

6. ☐ Microfiche Computer Program (Appendix)

7. Nucleotide and/or Amino Acid Sequence Submission (if applicable, all necessary)

a. ☐ Computer Readable Copy

b. ☐ Paper Copy (identical to computer copy)

c. ☐ Statement verifying identity of above copies

ACCOMPANYING APPLICATION PARTS

8. ☒ Assignment Papers (cover sheet & document(s))

9. ☐ 37 CFR 3.73(b) Statement

(when there is an assignee) ☐ Power of Attorney

10. ☐ English Translation Document (if applicable)

11. ☐ Information Disclosure Statement

(IDS)/PTO-1449 ☐ Copies of IDS Citations

12. ☐ Preliminary Amendment

13. ☒ Return Receipt Postcard (MPEP 503)

(Should be specifically itemized)

14. ☐ Certified Copy of Priority Document(s)
(if foreign priority is claimed)

15. ☐ Other:

16. ☐ If a CONTINUING APPLICATION, check appropriate box and supply the requisite information:

Amend the specification by inserting before the first line: -- This is a ☐ Continuation ☐ Divisional

☐ Continuation-in-Part (CIP) of prior application No.: , filed . --

17. For this divisional application, please cancel original Claims of the prior application before calculating the filing fee.

18. CORRESPONDENCE ADDRESS

☐ Customer Number or Bar Code Label

or ☒ Correspondence Address below

Name: Audley A. Ciamporcero, Jr., Esq.

Address: Johnson & Johnson

One Johnson & Johnson Plaza

New Brunswick, NJ 08933-7003 USA

19. TELEPHONE CONTACT

Please direct all telephone calls or telefaxes to Hal Brent Woodrow at:

Telephone: (732) 524-2976 Fax: (732) 524-2808

19. SIGNATURE OF APPLICANT, ATTORNEY, OR AGENT REQUIRED

NAME

Hal Brent Woodrow

Reg. No. 32501

SIGNATURE

Hal Brent Woodrow

DATE

June 26, 2000

06/26/00

06-27-00

A

jc057 U.S. PTO
09/603338



IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicant: Han-Cheng Zhang, Bruce E. Maryanoff, Anjali Pandey
and Robert M. Scarborough

For : NOVEL INDAZOLE PEPTIDOMIMETICS AS THROMBIN RECEPTOR
ANTAGONISTS

Express Mail Certificate

"Express Mail" mailing number: EL327264640US

Date of Deposit: June 26, 2000

I hereby certify that this complete application, including utility patent application and fee transmittal sheets, specification pages, claims, executed Declaration and Power of Attorney, and executed Assignment + cover sheet , is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 CFR 1.10 on the date indicated above and is addressed to the Assistant Commissioner for Patents, Washington, D.C. 20231.

Hal Brent Woodrow

(Typed or printed name of person mailing paper or fee)

Hal Brent Woodrow

(Signature of person mailing paper or fee)

NOVEL INDAZOLE PEPTIDOMIMETICS AS THROMBIN RECEPTOR ANTAGONISTS

5 FIELD OF THE INVENTION

This invention relates to certain novel thrombin receptor antagonists, their synthesis and their use for the treatment of diseases associated with thrombosis, restenosis, hypertension, heart failure, arrhythmia, inflammation,
10 angina, stroke, atherosclerosis, ischemic conditions, Angiogenesis related disorders, cancer, and neurodegenerative disorders.

BACKGROUND OF THE INVENTION

15 Thrombin is an important serine protease in hemostasis and thrombosis. One of the key actions of thrombin is cellular modulation via receptor activation. A functional human thrombin receptor (PAR-1), cloned by Coughlin in 1991 (T.-K. Vu, *Cell* **1991**, 64, 1057), was found to be a member of the G-protein coupled receptor (GPCR) superfamily. The receptor activation
20 putatively occurs by N-terminal recognition and proteolytic cleavage at the Arg-41/Ser-42 peptide bond to reveal a truncated N-terminus. This new receptor sequence, which has an SFLLRN (Ser-Phe-Leu-Leu-Arg-Asn) N-terminus acting as a tethered ligand to recognize a site on the receptor, can trigger activation and signal transduction leading to platelet aggregation. Since
25 1991, three other protease-activated receptors with extensive homology to the thrombin receptor, "PAR-2" (S. Nystedt, *Proc. Natl. Acad. Sci USA* **1994**, 91, 9208), "PAR-3" (H. Ishihara, *Nature* **1997**, 386, 502), and "PAR-4" (W.-F. Xu, *Proc. Natl. Acad. Sci USA* **1998**, 95, 6642), have been cloned. Thrombin receptor (PAR-1) specific antibody-induced blockade of the platelet thrombin
30 receptor has shown efficacy against arterial thrombosis in vivo (J. J. Cook *Circulation* **1995**, 91, 2961). Hence, antagonists of the thrombin receptor (PAR-1) are useful to block these protease-activated receptors and, as such, may be used to treat platelet mediated thrombotic disorders such as

ORT-1238


myocardial infarction, stroke, restenosis, angina, atherosclerosis, and ischemic conditions.

The thrombin receptor (PAR-1) has also been identified on other cell types: endothelial, fibroblast, renal, osteosarcoma, smooth muscle, myocytes, tumor, and neuronal/glia. Thrombin activation of endothelial cells upregulates P-selectin to induce polymorphonuclear leukocyte adhesion - an inflammatory response of the vessel wall (Y. Sugama, *J. Cell Biol.* 1992, 119, 935). In fibroblasts, thrombin receptor (PAR-1) activation induces proliferation and transmission of mitogenic signals (D. T. Hung, *J. Cell Biol.* **1992**, 116, 827). Thrombin has been implicated in osteoblast proliferation through its activation of osteoblast cells (D. N. Tatakis, *Biochem. Biophys. Res. Commun.* **1991**, 174, 181). Thrombin has been implicated in the regulation and retraction of neurons (K. Jalink, *J. Cell. Biol.* **1992**, 118, 411). Therefore, in this context, the antagonist compounds of this invention may also be useful against inflammation, osteoporosis, Angiogenesis related disorders, cancer, neurodegenerative disorders, hypertension, heart failure, arrhythmia, glomerulonephritis.

The compounds of the present invention are a structurally novel class of indazole peptidomimetics represented by the general formula (I) below.

SUMMARY OF THE INVENTION

The present invention is directed to structurally novel compounds represented by the following general formula (I):



ORT-1238

(I)

wherein

A₁ and A₂ are each independently a D- or L-amino acid selected from the group consisting of alanine, β-alanine, arginine, homoarginine, cyclohexylalanine, citrulline, cysteine (optionally substituted with C₁-C₄ alkyl, aryl, or arC₁-C₄ alkyl), 2,4-diaminobutyric acid (optionally substituted with acyl, C₁-C₄ alkyl, aroyl, amidino, or MeC(NH)-), 2,3 diaminopropionic acid (optionally substituted with acyl, C₁-C₄ alkyl, aroyl, amidino, or MeC(NH)-), glutamine, glycine, indanylglycine, lysine (optionally substituted with acyl, C₁-C₄ alkyl, aroyl, MeC(NH)-), valine, methionine, proline, serine (optionally substituted with C₁-C₄ alkyl, aryl, or arC₁-C₄ alkyl), homoserine (optionally substituted with C₁-C₄ alkyl, aryl, or arC₁-C₄ alkyl), tetrahydroisoquinoline-3-COOH, threonine (optionally substituted with C₁-C₄ alkyl, aryl, or arC₁-C₄ alkyl), ornithine (optionally substituted with acyl, C₁-C₄ alkyl, aroyl, MeC(NH)-), and an unsubstituted or substituted aromatic amino acid selected from the group consisting of phenylalanine, heteroarylalanine, naphthylalanine, homophenylalanine, histidine, tryptophan, tyrosine, arylglycine, heteroarylglycine, aryl-β-alanine, and heteroaryl-β-alanine wherein the substituents on the aromatic amino acid are independently selected from one or more of halogen, C₁-C₄ alkyl, C₁-C₄ alkoxy, hydroxy, C₁-C₄ alkoxycarbonyl, amino, amidino, guanidino, fluorinated C₁-C₄ alkyl, fluorinated C₁-C₄ alkoxy, C₁-C₄ alkylsulfonyl, C₁-C₄ alkylcarbonyl, cyano, aryl, heteroaryl, arC₁-C₄ alkyl, C₂-C₄ alkenyl, alkynyl, or nitro;

R₁ is selected from amino, C₁-C₈ alkylamino, C₁-C₈ dialkylamino, arylamino, arC₁-C₈ alkylamino, C₃-C₈ cycloalkylamino, heteroalkylC₁-C₈ alkylamino, heteroalkylC₁-C₈ alkyl-N-methylamino, C₁-C₈ dialkylaminoC₁-C₈ alkylamino, -N(C₁-C₈alkyl)-C₁-C₈ alkyl-N(C₁-C₈alkyl)₂, N(C₁-C₈alkyl)(C₁-C₈alkenyl), -N(C₁-C₈alkyl)(C₃-C₈cycloalkyl), heteroalkyl or substituted heteroalkyl wherein the substituent on the heteroalkyl is selected from oxo, amino, C₁-C₈ alkoxyC₁-C₈ alkyl, C₁-C₈ alkylamino or C₁-C₈ dialkylamino;

ORT-1238

Preferably, R_1 is selected from amino, C_1 - C_6 alkylamino, C_1 - C_6 dialkylamino, arylamino, ar C_1 - C_6 alkylamino, heteroalkyl C_1 - C_6 alkylamino, -N(C_1 - C_6 alkyl)- C_1 - C_6 alkyl-N(C_1 - C_6 alkyl)₂, heteroalkyl or substituted heteroalkyl wherein the substituent on the heteroalkyl is selected from oxo, amino, C_1 - C_6 alkoxy C_1 - C_6 alkyl, C_1 - C_6 alkylamino or C_1 - C_6 dialkylamino;

R_2 and R_3 are each independently selected from hydrogen, C_1 - C_8 alkyl, C_3 - C_8 cycloalkyl, C_3 - C_8 cycloalkyl C_1 - C_8 alkyl, aryl, heteroalkyl, substituted heteroalkyl (wherein the substituent on the heteroalkyl is one or more substituents independently selected from C_1 - C_8 alkoxycarbonyl, C_1 - C_8 alkyl, or C_1 - C_4 alkylcarbonyl), heteroalkyl C_1 - C_8 alkyl, indanyl, acetamidino C_1 - C_8 alkyl, amino C_1 - C_8 alkyl, C_1 - C_8 alkylamino C_1 - C_8 alkyl, C_1 - C_8 dialkylamino C_1 - C_8 alkyl, unsubstituted or substituted heteroaryl C_1 - C_8 alkyl or unsubstituted or substituted ar C_1 - C_8 alkyl, wherein the substituent on the aralkyl or heteroarylalkyl group is one or more substituents independently selected from halogen, nitro, amino, C_1 - C_8 alkyl, C_1 - C_8 alkoxy, hydroxy, cyano, C_1 - C_4 alkylcarbonyl, C_1 - C_8 alkoxycarbonyl, hydroxy C_1 - C_8 alkyl or aminosulfonyl; or

R_2 and R_3 together with the nitrogen to which they are attached, alternatively form an unsubstituted or substituted heteroalkyl group selected from piperidinyl, piperazinyl, morpholinyl or pyrrolidinyl, wherein the substituent is one or more substituents independently selected from C_1 - C_8 alkyl, C_1 - C_8 alkoxycarbonyl or C_1 - C_4 alkylcarbonyl;

Preferably, R_2 is selected from hydrogen or C_1 - C_6 alkyl; and

R_3 is selected from C_1 - C_8 alkyl, C_3 - C_6 cycloalkyl, C_3 - C_6 cycloalkyl C_1 - C_6 alkyl, aryl, heteroaryl C_1 - C_6 alkyl, substituted heteroaryl C_1 - C_6 alkyl wherein the substituent is C_1 - C_4 alkyl, heteroalkyl, heteroalkyl C_1 - C_6 alkyl, indanyl, acetamidino C_1 - C_6 alkyl, amino C_1 - C_6 alkyl, C_1 - C_6 alkylamino C_1 - C_6 alkyl, C_1 - C_6 dialkylamino C_1 - C_6 alkyl, ar C_1 - C_6 alkyl, substituted ar C_1 - C_6 alkyl wherein the substituent on the aralkyl group is one to five substituents independently selected from halogen, nitro, amino, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, C_1 - C_6 alkoxycarbonyl, hydroxyalkyl or aminosulfonyl; or

R_2 and R_3 , together with the nitrogen to which they are attached, alternatively form an unsubstituted or substituted heteroalkyl group selected

ORT-1238

from piperidinyl, piperazinyl or pyrrolidinyl, wherein the substituent is independently one or two substituents selected from C₁-C₆ alkyl;

R₄ is selected from unsubstituted or substituted aryl, arC₁-C₈ alkyl, C₃-C₈ cycloalkyl, or heteroaryl, where the substituents on the aryl, arC₁-C₈ alkyl, cycloalkyl or heteroaryl group are independently selected from one or more of halogen, nitro, amino, cyano, hydroxyalkyl, C₁-C₈ alkyl, C₁-C₈ alkoxy, hydroxy, C₁-C₄ alkylcarbonyl, C₁-C₈ alkoxycarbonyl, fluorinated C₁-C₄ alkyl, fluorinated C₁-C₄ alkoxy or C₁-C₄ alkylsulfonyl;

Preferably, R₄ is selected from unsubstituted or substituted aryl, arC₁-C₆ alkyl, C₃-C₆ cycloalkyl or heteroaryl, where the substituents on the aryl, aralkyl, cycloalkyl or heteroaryl group are independently selected from one to three substituents selected from halogen, cyano, C₁-C₄ alkyl, C₁-C₄ alkoxy, C₁-C₄ alkoxycarbonyl, fluorinated C₁-C₄ alkyl, fluorinated C₁-C₄ alkoxy or C₁-C₄ alkylsulfonyl;

R₅ is selected from hydrogen or C₁-C₈ alkyl; preferably, R₅ is hydrogen

X is oxygen or sulfur; preferably, X is oxygen;

m is an integer selected from 0, 1, 2 or 3;

n is an integer selected from 1 or 2;

p is an integer selected from 0 or 1; preferably, p is 1;

and pharmaceutically acceptable salts thereof.

In a preferred embodiment of the present invention:

A₁ is an L-amino acid selected from the group consisting of alanine, arginine, cyclohexylalanine, glycine, proline, tetrahydroisoquinoline-3-COOH, and an unsubstituted or substituted aromatic amino acid selected from the group consisting of phenylalanine, naphthylalanine, homophenylalanine, and O-methyl tyrosine, wherein the substituents on the aromatic amino acid are independently selected from one to five of (preferably, one to three of) halogen, C₁-C₄ alkyl, C₁-C₄ alkoxy, hydroxy, C₁-C₄ alkoxycarbonyl, amino,

ORT-1238

amidino, guanidino, fluorinated C₁-C₄ alkyl, fluorinated C₁-C₄ alkoxy, C₁-C₄ alkylsulfonyl, C₁-C₄ alkylcarbonyl, cyano, aryl, heteroaryl, arC₁-C₄ alkyl, C₂-C₄ alkenyl, alkynyl, or nitro;

- 5 A₂ is an L-amino acid selected from the group consisting of alanine, β-alanine, arginine, citrulline, cysteine (optionally substituted with C₁-C₄ alkyl, aryl, or arC₁-C₄ alkyl), 2,4-diaminobutyric acid (optionally substituted with acyl, C₁-C₄ alkyl, aroyl, amidino, or MeC(NH)-), 2,3- diaminopropionic acid (optionally substituted with acyl, C₁-C₄ alkyl, aroyl, amidino, or MeC(NH)-),
- 10 glutamine, glycine, lysine (optionally substituted with acyl, C₁-C₄ alkyl, aroyl, MeC(NH)-), valine, methionine, serine (optionally substituted with C₁-C₄ alkyl, aryl, or arC₁-C₄ alkyl), homoserine (optionally substituted with C₁-C₄ alkyl, aryl, or arC₁-C₄ alkyl), threonine (optionally substituted with C₁-C₄ alkyl, aryl, or arC₁-C₄ alkyl), ornithine (optionally substituted with acyl, C₁-C₄ alkyl, aroyl,
- 15 MeC(NH)-), and an unsubstituted or substituted aromatic amino acid selected from the group consisting of phenylalanine, heteroarylalanine, and histidine, wherein the substituents on the aromatic amino acid are independently selected from one to five of (preferably, one to three of) halogen, C₁-C₄ alkyl, C₁-C₄ alkoxy, hydroxy, C₁-C₄ alkoxycarbonyl, amino, amidino, guanidino,
- 20 fluorinated C₁-C₄ alkyl, fluorinated C₁-C₄ alkoxy, C₁-C₄ alkylsulfonyl, C₁-C₄ alkylcarbonyl, cyano, aryl, heteroaryl, arC₁-C₄ alkyl, C₂-C₄ alkenyl, alkynyl, or nitro;

- R₂ is selected from hydrogen or C₁-C₄ alkyl;
- 25 m and n are both 1;

and all other variables are as defined previously;
and pharmaceutically acceptable salts thereof.

- 30 In a class of the invention:


A₁ is an L-amino acid selected from the group consisting of alanine, arginine, cyclohexylalanine, glycine, proline, and an unsubstituted or substituted aromatic amino acid selected from the group consisting of

ORT-1238


phenylalanine, naphthylalanine, homophenylalanine, and O-methyl tyrosine, wherein the substituents on the aromatic amino acid are independently one to two substituents selected from halogen, C₁-C₄ alkyl, C₁-C₄ alkoxy, hydroxy, C₁-C₄ alkoxy carbonyl, amino, amidino, guanidino, fluorinated C₁-C₄ alkyl, fluorinated C₁-C₄ alkoxy, C₁-C₄ alkylsulfonyl, C₁-C₄ alkylcarbonyl, cyano, aryl, heteroaryl, arC₁-C₄ alkyl, C₂-C₄ alkenyl, alkynyl, or nitro;

A₂ is an L-amino acid selected from the group consisting of alanine, β-alanine, arginine, citrulline, cysteine (optionally substituted with C₁-C₄ alkyl, aryl, or arC₁-C₄ alkyl), 2,4-diaminobutyric acid (optionally substituted with acyl, C₁-C₄ alkyl, aroyl, amidino, or MeC(NH)-), 2,3-diaminopropionic acid (optionally substituted with acyl, C₁-C₄ alkyl, aroyl, amidino, or MeC(NH)-), glutamine, glycine, lysine (optionally substituted with acyl, C₁-C₄ alkyl, aroyl, MeC(NH)-), valine, methionine, serine (optionally substituted with C₁-C₄ alkyl, aryl, or arC₁-C₄ alkyl), homoserine (optionally substituted with C₁-C₄ alkyl, aryl, or arC₁-C₄ alkyl), threonine (optionally substituted with C₁-C₄ alkyl, aryl, or arC₁-C₄ alkyl), ornithine (optionally substituted with acyl, C₁-C₄ alkyl, aroyl, MeC(NH)-), and an unsubstituted or substituted aromatic amino acid selected from the group consisting of phenylalanine, heteroarylalanine, and histidine, wherein the substituents on the aromatic amino acid are independently one to two substituents selected from halogen, C₁-C₄ alkyl, C₁-C₄ alkoxy, hydroxy, C₁-C₄ alkoxy carbonyl, amino, amidino, guanidino, fluorinated C₁-C₄ alkyl, fluorinated C₁-C₄ alkoxy, C₁-C₄ alkylsulfonyl, C₁-C₄ alkylcarbonyl, cyano, aryl, heteroaryl, arC₁-C₄ alkyl, C₂-C₄ alkenyl, alkynyl, or nitro;

R₁ is selected from diethylamino, di-(*n*-propyl)amino,



ORT-1238

Preferably, R₁ is:  ;

R₂ is selected from hydrogen, methyl or ethyl;

5 R₃ is selected from 2-indanyl, phenyl, cyclohexylmethyl, cyclopentyl, pyridylmethyl, furanylmethyl, 2-(4-methyl-furanyl)methyl, thienylmethyl, diphenylmethyl, 4-imidazolylethyl, 2-(4-N-methyl)imidazolylethyl, *n*-octyl, phenyl-*n*-propyl, aminoethyl, aminopropyl, amino-*n*-pentyl, dimethylaminoethyl, 4-aminophenylsulfonylaminoethyl, acetamidineylethyl, 2-N-pyrrolidinylethyl, 10 N-ethoxycarbonylpiperidinyl, unsubstituted or substituted phenylethyl or unsubstituted or substituted benzyl wherein the substituents on the phenylethyl or benzyl are independently one or two substituents selected from methyl, fluorine, chlorine, nitro, methoxy, methoxycarbonyl or hydroxymethyl; or

15 R₂ and R₃, together with the nitrogen to which they are attached, form a heteroalkyl group selected from piperidinyl, or 4-(N-methyl)piperazinyl;

R₄ is selected from cyclohexyl, 2-naphthyl, phenylethyl, 4-fluorophenylethyl, or unsubstituted or substituted phenyl, where the 20 substituents on the phenyl are independently selected from one to two substituents selected from fluorine, chlorine, iodine, methyl cyano, or trifluoromethyl;

25 Preferably, R₄ is 2,6-dichlorophenyl or 2-methylphenyl;

all other variables are as defined previously;
and pharmaceutically acceptable salts thereof.

In a subclass of the invention,

30 A₁ is selected from 3,4-Difluorophenylalanine or 4-Chlorophenylalanine;

A₂ is selected from 2,4-Diaminobutyric acid or 4-Pyridylalanine;

R₂ is hydrogen;

R₃ is selected from benzyl or 2-aminoethyl;

35

ORT-1238

all other variables are as defined previously;
and pharmaceutically acceptable salts thereof.


Illustrative of the invention is a pharmaceutical composition comprising a
5 pharmaceutically acceptable carrier and any of the compounds described
above. Illustrating the invention is a pharmaceutical composition made by
mixing any of the compounds described above and a pharmaceutically
acceptable carrier. An illustration of the invention is a process for making a
pharmaceutical composition comprising mixing any of the compounds
10 described above and a pharmaceutically acceptable carrier.

An example of the invention is a method of treating a disorder
(preferably, a platelet-mediated thrombotic disorder) selected from arterial
and/or venous thrombosis, acute myocardial infarction, reocclusion following
15 thrombolytic therapy and/or angioplasty, inflammation, unstable angina, stroke,
restenosis, atherosclerosis, ischemic conditions, hypertension, heart failure,
arrhythmia, glomerulonephritis, osteoporosis, Angiogenesis related disorders,
cancer, neurodegenerative disorders and a variety of vaso-occlusive disorders
in a subject in need thereof comprising administering to the subject a
20 therapeutically effective amount of any of the compounds or pharmaceutical
compositions described above. In a preferred embodiment, the therapeutically
effective amount of the compound is from about 0.1 mg/kg/day to about 300
mg/kg/day.


Also included in the invention is the use of any of the compounds
described above for the preparation of a medicament for a disorder (preferably,
a platelet-mediated thrombotic disorder) selected from arterial and/or venous
thrombosis, acute myocardial infarction, reocclusion following thrombolytic
therapy and/or angioplasty, inflammation, unstable angina, stroke, restenosis,
25 atherosclerosis, ischemic conditions, hypertension, heart failure, arrhythmia,
glomerulonephritis, osteoporosis, Angiogenesis related disorders, cancer,
neurodegenerative disorders or a variety of vaso-occlusive disorders in a
subject in need thereof.

DETAILED DESCRIPTION OF THE INVENTION

More particularly, the present invention is directed to compounds of the following formula (I):



wherein A_1 , A_2 , R_1 , R_2 , R_3 , R_4 , R_5 , X , m , n and p are as previously defined. In a particularly preferred embodiment, the compounds have the formula




The compounds of the present invention are thrombin receptor antagonists and as such are useful in treating thrombosis, restenosis, hypertension, heart failure, arrhythmia, myocardial infarction, glomerulonephritis, reocclusion following thrombolytic therapy, reocclusion following angioplasty, inflammation, angina, stroke, atherosclerosis, ischemic conditions, a vaso-occlusive disorder, neurodegenerative disorders, Angiogenesis related disorders and cancer. These compounds are also useful as antithrombotics in conjunction with fibrinolytic therapy (e.g., t-PA or streptokinase).

In the compounds of formula (I), the amino acid residues comprising the A_1 and A_2 substituents are attached to the adjacent moiety according to standard nomenclature so that the amino-terminus (N-terminus) of the amino acid is drawn on the left and the carboxy-terminus of the amino acid is drawn on the right. So, for example, in Compound 1, where A_1 is 3,4-difluorophenylalanine and A_2 is Dbu (2,4-Diaminobutyric acid), the N-terminus of the 3,4-difluorophenylalanine (A_1) is attached to the carbonyl group and the carboxy-terminus of the 3,4-difluorophenylalanine (A_1) is attached to the N-terminus of the A_2 substituent (Dbu), similarly, the N-terminus of the Dbu (A_2) is attached to the carboxy-terminus of the A_1 substituent and the carboxy-terminus of the Dbu (A_2) is attached to the $N-R_2R_3$ group.

When a particular group is "substituted" (e.g., Phe, aryl, heteroalkyl, heteroaryl), that group may have one or more substituents, preferably from one to five substituents, more preferably from one to three substituents, most preferably from one to two substituents, independently selected from the list of substituents.

Under standard nomenclature used throughout this disclosure, the terminal portion of the designated side chain is described first, followed by the adjacent functionality toward the point of attachment. Thus, for example, a "phenylC₁-C₆ alkylamidoC₁-C₆alkyl" substituent refers to a group of the formula



The compounds of the present invention may also be present in the form of a pharmaceutically acceptable salt. The pharmaceutically acceptable salt generally takes a form in which the basic nitrogen is protonated with an inorganic or organic acid. Representative organic or inorganic acids include hydrochloric, hydrobromic, hydriodic, perchloric, sulfuric, nitric, phosphoric, acetic, propionic, glycolic, lactic, succinic, maleic, fumaric, malic, tartaric, citric, benzoic, mandelic, methanesulfonic, hydroxyethanesulfonic, benzenesulfonic,

ORT-1238

oxalic, pamoic, 2-naphthalenesulfonic, p-toluenesulfonic, cyclohexanesulfamic, salicylic, saccharinic or trifluoroacetic.

Where the compounds according to this invention have at least one
5 chiral center, they may accordingly exist as enantiomers. Where the compounds possess two or more chiral centers, they may additionally exist as diastereomers. It is to be understood that all such isomers and mixtures thereof are encompassed within the scope of the present invention. Furthermore, some of the crystalline forms for the compounds may exist as
10 polymorphs and as such are intended to be included in the present invention. In addition, some of the compounds may form solvates with water (i.e., hydrates) or common organic solvents, and such solvates are also intended to be encompassed within the scope of this invention.

15 The term "subject" as used herein, refers to an animal, preferably a mammal, most preferably a human, who has been the object of treatment, observation or experiment.

The term "therapeutically effective amount" as used herein, means that
20 amount of active compound or pharmaceutical agent that elicits the biological or medicinal response in a tissue system, animal or human that is being sought by a researcher, veterinarian, medical doctor or other clinician, which includes alleviation of the symptoms of the disease or disorder being treated.

25 As used herein, unless otherwise noted alkyl and alkoxy whether used alone or as part of a substituent group, include straight and branched chains having 1 to 8 carbon atoms, or any number within this range. For example, alkyl radicals include methyl, ethyl, propyl, isopropyl, *n*-butyl, isobutyl, *sec*-butyl, *t*-butyl, *n*-pentyl, 3-(2-methyl)butyl, 2-pentyl, 2-methylbutyl, neopentyl, *n*-
30 hexyl, 2-hexyl and 2-methylpentyl. Alkoxy radicals are oxygen ethers formed from the previously described straight or branched chain alkyl groups. Cycloalkyl groups contain 3 to 8 ring carbons and preferably 5 to 7 carbons. Similarly, alkenyl and alkynyl groups include straight and branched chain

ORT-1238

alkenes and alkynes having 1 to 8 carbon atoms, or any number within this range.

5 The term "aryl" as used herein refers to an unsubstituted or substituted aromatic group such as phenyl and naphthyl. The term "aroyl" refers to the group -C(O)-aryl .

10 The term "heteroalkyl" as used herein represents an unsubstituted or substituted stable three to seven membered monocyclic saturated ring system which consists of carbon atoms and from one to three heteroatoms selected from N, O or S, and wherein the nitrogen or sulfur heteroatoms may optionally be oxidized, and the nitrogen heteroatom may optionally be quaternized. The heteroalkyl group may be attached at any heteroatom or carbon atom which
15 results in the creation of a stable structure. Examples of such heteroalkyl groups include, but are not limited to azetidiny, piperidiny, pyrrolidiny, piperaziny, oxopiperaziny, oxopiperidiny, oxoazepiny, azepiny, tetrahydrofurany, dioxolany, tetrahydroimidazoly, tetrahydrothiazoly, tetrahydrooxazoly, tetrahydropyrany, morpholiny, thiomorpholiny, thiamorpholiny sulfoxide, thiamorpholiny sulfone and oxadiazoly. Preferred
20 heteroalkyl groups include pyrrolidiny, piperidiny, piperaziny, morpholiny, azetidiny and tetrahydrothiazoly.

25 The term "heteroaryl" as used herein represents an unsubstituted or substituted stable five or six membered monocyclic aromatic ring system or an unsubstituted or substituted nine or ten membered benzo-fused heteroaromatic ring system or bicyclic heteroaromatic ring system which consists of carbon atoms and from one to four heteroatoms selected from N, O or S, and wherein the nitrogen or sulfur heteroatoms may optionally be oxidized, and the nitrogen
30 heteroatom may optionally be quaternized. The heteroaryl group may be attached at any heteroatom or carbon atom that results in the creation of a stable structure. Examples of heteroaryl groups include, but are not limited to

ORT-1238

pyridyl, pyridazinyl, thienyl, furanyl, imidazolyl, isoxazolyl, oxazolyl, pyrazolyl, pyrrolyl, thiazolyl, thiadiazolyl, triazolyl, benzimidazolyl, benzofuranyl, benzothienyl, benzisoxazolyl, benzoxazolyl, benzopyrazolyl, indolyl, benzothiazolyl, benzothiadiazolyl, benzotriazolyl adeninyl or quinolinyl.

- 5 Preferred heteroaryl groups include pyridyl, pyrrolyl, pyrazinyl, thiadiazolyl, pyrazolyl, thienyl, triazolyl and quinolinyl.

The term "aralkyl" means an alkyl group substituted with one, two or three aryl groups (e.g., benzyl, phenylethyl, diphenylmethyl, triphenylmethyl).

- 10 Similarly, the term "aralkoxy" indicates an alkoxy group substituted with an aryl group (e.g., benzyloxy). The term aminoalkyl refers to an alkyl group substituted with an amino group (*i.e.*, -alkyl-NH₂). The term "alkylamino" refers to an amino group substituted with an alkyl group (*i.e.*, -NH-alkyl). The term "dialkylamino" refers to an amino group which is disubstituted with alkyl groups
15 wherein the alkyl groups can be the same or different (*i.e.*, -N-[alkyl]₂).

The term "acyl" as used herein means an organic radical having 1 to 6 carbon atoms (branched or straight chain) derived from an organic acid by removal of the hydroxyl group.

20

The term "oxo" refers to the group =O.

The term "carbonyl" refers to the group C(O).

25

The term "halogen" shall include iodine, bromine, chlorine and fluorine.

Whenever the term "alkyl" or "aryl" or either of their prefix roots appear in a name of a substituent (e.g., aralkyl, dialkylamino) it shall be interpreted as including those limitations given above for "alkyl" and "aryl." Designated
30 numbers of carbon atoms (e.g., C₁-C₆) shall refer independently to the number of carbon atoms in an alkyl or cycloalkyl moiety or to the alkyl portion of a larger substituent in which alkyl appears as its prefix root.


As used herein, the term "phosgene equivalent" represents the class of carbonic acid derivatives which include 4-nitrophenyl chloroformate, phosgene or "COCl₂," phenyl chloroformate, triphosgene or "(CCl₃O)₂CO," carbonyldiimidazole, diethyl carbonate or diphenyl carbonate.

It is intended that the definition of any substituent or variable at a particular location in a molecule be independent of its definitions elsewhere in that molecule. It is understood that substituents and substitution patterns on the compounds of this invention can be selected by one of ordinary skill in the art to provide compounds that are chemically stable and that can be readily synthesized by techniques known in the art as well as those methods set forth herein.

As used herein, the term "composition" is intended to encompass a product comprising the specified ingredients in the specified amounts, as well as any product which results, directly or indirectly, from combinations of the specified ingredients in the specified amounts. Accordingly, pharmaceutical compositions containing the compounds of the present invention as the active ingredient as well as methods of preparing the instant compounds are also part of the present invention.

Particularly preferred compounds of the present invention and their biological data are shown in Table 1, as follows; the amino acids bear the "L" absolute configuration unless denoted otherwise. Table 1 contains IC₅₀ values (μM) of the compounds in a thrombin receptor binding assay, and IC₅₀ values (μM) against platelet aggregation stimulated by thrombin.

Table 1. Indazole Peptidomimetics As
Thrombin Receptor (PAR-1) Antagonists



5

| Comp | A ₁ | A ₂ | R ₂ R ₃ N | IC ₅₀ (μM) | |
|------|--------------------------|-----------------------|--|---------------------------|-----------------------------|
| | | | | Thr GFP Aggr ^a | Thr Recptr Bdg ^b |
| 1 | 3,4-DiF-Phe ^c | Dbu ^d | PhCH ₂ NH | 0.31 | 0.04 |
| 2 | 4-Cl-Phe | Dbu | PhCH ₂ NH | 0.26 | 20 |
| 3 | 3,4-DiF-Phe | 4-Pyrala ^e | H ₂ NCH ₂ CH ₂ NH | 0.50 | 0.03 |
| 4 | 3,4-DiF-Phe | Dbu | R-PhCH(Me)NH | 0.32 | 0.15 |
| 5 | 3,4-DiF-Phe | Dbu | S-PhCH(CH ₂ OH)NH | 0.66 | 0.32 |
| 6 | 4-Cl-Phe | 2-Thiala ^f | H ₂ NCH ₂ CH ₂ NH | 0.30 | 5.8 |

^a Thrombin-induced gel-filtered platelet aggregation assay.

^b Thrombin receptor (PAR-1) binding assay.

^c 3,4-Difluorophenylalanine. ^d 2,4-Diaminobutyric acid. ^e 4-Pyridylalanine.

^f 2-Thienylalanine.

10

The antagonists of the present invention may be prepared via a convergent solution-phase synthesis by coupling an aminoindazole intermediate **AAG4** with a dipeptide amine **AAG6** via a urea linkage as described in the general Scheme AAGeneric. The appropriately nitro substituted indole **AAG1** (Scheme AAGeneric) was treated with aqueous NaNO₂ under acidic conditions (pH from about pH 1 to about pH 2) to give (via nitrosation, G. Buchi, *J. Am. Chem. Soc.* **1986**, *108*, 4115) 3-indazolecarboxaldehyde **AAG2**. Reductive amination of **AAG2** with an amine such as pyrrolidine and a reducing agent such as sodium triacetoxyborohydride afforded **AAG3**. Alkylation of **AAG3** with a substituted alkyl or heteroaryl alkyl halide and a base such as potassium hydroxide in an

15

20

ORT-1238

aprotic solvent such as THF to give an intermediate, which was reduced in a classical manner with, for example, iron and acetic acid or with a newer method such as dimethyl hydrazine and iron to give aminoindazole intermediate **AAG4**.

5

Dipeptide amine **AAG6** can be synthesized from the corresponding protected amino acids using standard peptide coupling conditions. Thus, an Fmoc protected amino-acid (A_2), **AAG5** (Scheme AAGeneric), was coupled to amine R_2R_3NH using a coupling agent such as dicyclohexylcarbodiimide (DCC) or diisopropylcarbodiimide (DIC) and 1-hydroxybenzotriazole (HOBT) in a dipolar aprotic solvent like DMF to give the amide, which was Fmoc deprotected with a dialkylamine in a dipolar aprotic solvent such as diethylamine in acetonitrile. The resulting amine was coupled to the second Fmoc protected amino-acid (A_1) in the same way with a coupling agent such as DIC and HOBT in a dipolar aprotic solvent like DMF to give the dipeptide, which was Fmoc deprotected as above with a dialkylamine in a dipolar aprotic solvent like acetonitrile to afford dipeptide amine **AAG6**.


10

15

20

Aminoindazole intermediate **AAG2** was then treated with a phosgene equivalent such as 4-nitrophenyl chloroformate or triphosgene and a base like diisopropylethylamine in a solvent such as dichloromethane, and to this was then added dipeptide amine **AAG6** to give an urea. Removal of the protecting group, if necessary, such as Boc group with an acid such as trifluoroacetic acid from the side chain of dipeptide afforded final targets **AAG7**.


SCHEME AAGeneric



ORT-1238

As a typical example of this convergent solution-phase method, synthesis of compound **1** was presented in Scheme AA. Thus, treatment of 6-nitroindole **AA1** with aqueous NaNO_2 under acidic condition (pH from about pH 1 to about pH 2) afforded 3-indazolecarboxaldehyde (**AA2**). Reductive amination of **AA2** with pyrrolidine/ $\text{NaB}(\text{OAc})_3\text{H}$ was followed by alkylation with 2,6-diCl-Bn-Br and nitro reduction with $\text{Me}_2\text{NNH}_2/\text{FeCl}_3$ to provide aminoindazole intermediate **AA4**. Coupling of N- α -Fmoc-N- γ -Boc-diaminobutyric acid (**AA5**) with benzyl amine in the presence of DCC and HOBt was followed by de-protection of Fmoc group with diethylamine. The resulting intermediate was coupled with Fmoc-3,4-diF-Phe-OH using DIC/HOBt and treated with diethylamine to give dipeptide amine **AA6**. Urea formation between dipeptide amine **AA6** and 6-aminoindazole **AA4** in the presence of 4-nitrophenylchloroformate was followed by de-protection of Boc group with TFA to afford target compound **1**.

SCHEME AA




ORT-1238


Alternatively, the antagonists of the present invention may also be prepared via solid-phase methods as represented by the synthesis of **2** and **3** (Schemes AB and Scheme AC, respectively). In Scheme AB, N- α -Fmoc-N- γ -Boc-2,4-diaminobutyric acid (**AB1**) was coupled with benzyl amine in the presence of DCC and HOBt. The resulting benzylamide was treated with TFA in DCM to give **AB2**, which was then loaded onto 2-Cl-trityl-Cl resin in the presence of DIEA to afford **AB3**. Deprotection of Fmoc group in **AB3** with piperidine was followed by coupling with Fmoc-4-Cl-Phe-OH in the presence of HBTU and HOBt. The resulting coupled product was deprotected again with piperidine to afford the resin-bound dipeptide amine **AB4**. Urea formation between **AB4** and aminoindazole intermediate **AA4** was accomplished by using 4-nitrophenylchloroformate to provide **AB5**, which was cleaved with TFA to afford target **2**.

Similarly, Scheme AC described a solid-phase synthesis of the antagonists having an amine group at carboxy-terminus of the A₂, such as **3** and **6**, by mono-attaching a di-amine, such as ethylenediamine, on 2-Cl-trityl-Cl resin followed by coupling with the protected amino acid A₂ and then A₁ to furnish the required resin-bound dipeptide amine such as **AC4**.

SCHEME AB



SCHEME AC



ORT-1238

The side-chain amine in antagonists such as **1** and **3** may be converted to other functional groups such as acetamidine and guanidine by using standard procedures. For example, the acetamidine and guanidine groups can be introduced by treating the side-chain amine with S-2-naphthylmethyl thioacetimidate hydrobromide and 2-methyl-2-thiopseudourea, respectively.

The thioureidoindoles [X = S, general formula (I)] may be prepared as described hereinafter. Aminoindazole substrate is reacted with thiocarbonyldiimidazole in a chlorinated solvent and then the imidazole by-product filtered from the solution. The solution then can be concentrated to afford the N-imidazolyl-N'-aminoindazolyl-thiourea. This intermediate is then reacted with a peptide amine in a polar, aprotic solvent with heating (80-100 degrees) to afford the N-peptido-N'-aminoindazolyl-thiourea product.

Amidoindazoles [p = 0, X = O, general formula (I)] may be prepared from a dipeptide amine **AAG6** (Scheme AAGeneric) and an indazole carboxylic acid intermediate by using standard coupling conditions such as DCC/HOBt. The required indazole carboxylic acid intermediates can be prepared from the appropriately indole carboxylic acid esters by using the same method as described for aminoindazole intermediate **AAG4** in Scheme AAGeneric.

Carbon-chain extension from n = 1 to n = 2 at the 3-position of the indazole [see general formula (I)] may be introduced in the intermediate **AAG2** (Scheme AAGeneric) via aldehyde-nitromethane condensation followed by reduction of the resulting α,β -unsaturated nitro compounds to saturated amine.

The utility of the compounds to treat PAR-1 mediated disorders (e.g., thrombotic disorders) can be determined according to the procedures described herein. The present invention therefore provides a method of treating PAR-1 mediated disorders (e.g., thrombotic disorders) in a subject in need thereof which comprises administering any of the compounds as defined

ORT-1238

herein in a quantity effective to treat PAR-1 mediated disorders. The compound may be administered to a patient by any conventional route of administration, including, but not limited to, intravenous, oral, subcutaneous, intramuscular, intradermal and parenteral.

5

The present invention also provides pharmaceutical compositions comprising one or more compounds of this invention in association with a pharmaceutically acceptable carrier.

10

To prepare the pharmaceutical compositions of this invention, one or more compounds of formula (I) or salt thereof of the invention as the active ingredient, is intimately admixed with a pharmaceutical carrier according to conventional pharmaceutical compounding techniques, which carrier may take a wide variety of forms depending on the form of preparation desired for

15

administration, e.g., oral or parenteral such as intramuscular. In preparing the compositions in oral dosage form, any of the usual pharmaceutical media may be employed. Thus, for liquid oral preparations, such as, for example,

20

suspensions, elixirs and solutions, suitable carriers and additives include water, glycols, oils, alcohols, flavoring agents, preservatives, coloring agents and the like; for solid oral preparations such as, for example, powders, capsules, caplets, gelcaps and tablets, suitable carriers and additives include starches, sugars, diluents, granulating agents, lubricants, binders, disintegrating agents and the like. Because of their ease in administration, tablets and capsules represent the most advantageous oral dosage unit form,

25

in which case solid pharmaceutical carriers are obviously employed. If desired, tablets may be sugar coated or enteric coated by standard techniques. For parenterals, the carrier will usually comprise sterile water, though other ingredients, for example, for purposes such as aiding solubility or for preservation, may be included. Injectable suspensions may also be prepared,

30

in which case appropriate liquid carriers, suspending agents and the like may be employed. The pharmaceutical compositions herein will contain, per dosage unit, e.g., tablet, capsule, powder, injection, teaspoonful and the like, an amount of the active ingredient necessary to deliver an effective dose as described above. The pharmaceutical compositions herein will contain, per

35

unit dosage unit, e.g., tablet, capsule, powder, injection, suppository, teaspoonful and the like, from about 0.03 mg/kg to about 100 mg/kg (preferred

ORT-1238

from about 0.1 mg/kg to about 30 mg/kg) of a compound of the present invention and may be given at a dosage from about 0.1 mg/kg/day to about 300 mg/kg/day (preferred from about 1 mg/kg/day to about 50 mg/kg/day). The dosages, however, may be varied depending upon the requirement of the patients, the severity of the condition being treated and the compound being employed. The use of either daily administration or post-periodic dosing may be employed.

Preferably these compositions are in unit dosage forms such as tablets, pills, capsules, powders, granules, sterile parenteral solutions or suspensions, metered aerosol or liquid sprays, drops, ampoules, autoinjector devices or suppositories for oral parenteral, intranasal, sublingual or rectal administration, or for administration by inhalation or insufflation. Alternatively, the composition may be presented in a form suitable for once-weekly or once-monthly administration; for example, an insoluble salt of the active compound, such as the decanoate salt, may be adapted to provide a depot preparation for intramuscular injection. For preparing solid compositions such as tablets, the principal active ingredient is mixed with a pharmaceutical carrier, e.g. conventional tableting ingredients such as corn starch, lactose, sucrose, sorbitol, talc, stearic acid, magnesium stearate, dicalcium phosphate or gums and other pharmaceutical diluents, e.g. water, to form a solid preformulation composition containing a homogeneous mixture of a compound of the present invention or a pharmaceutically acceptable salt thereof. When referring to these preformulation compositions as homogeneous, it is meant that the active ingredient is dispersed evenly throughout the composition so that the composition may be readily subdivided into equally effective dosage forms such as tablets, pills and capsules. This solid preformulation composition is then subdivided into unit dosage forms of the type described above containing from about 0.1 mg to about 500 mg of the active ingredient of the present invention. The tablets or pills of the novel composition can be coated or otherwise compounded to provide a dosage form affording the advantage of prolonged action. For example, the tablet or pill can comprise an inner dosage and an outer dosage component, the latter being in the form of an envelope over the former. The two components can be separated by an enteric layer

ORT-1238

which serves to resist disintegration in the stomach and permits the inner component to pass intact into the duodenum or to be delayed in release. A variety of material can be used for such enteric layers or coatings, such materials including a number of polymeric acids with such materials as shellac, 5 cetyl alcohol and cellulose acetate.

The liquid forms in which the novel compositions of the present invention may be incorporated for administration orally or by injection include aqueous solutions, suitably flavoured syrups, aqueous or oil suspensions and 10 flavoured emulsions with edible oils such as cottonseed oil, sesame oil, coconut oil or peanut oil, as well as elixirs and similar pharmaceutical vehicles. Suitable dispersing or suspending agents for aqueous suspensions, include synthetic and natural gums such as tragacanth, acacia, alginate, dextran, sodium carboxymethylcellulose, methylcellulose, polyvinyl-pyrrolidone or 15 gelatin.

Where the processes for the preparation of the compounds according to the invention give rise to mixture of stereoisomers, these isomers may be separated by conventional techniques such as preparative chromatography. 20 The compounds may be prepared in racemic form, or individual enantiomers may be prepared either by enantiospecific synthesis or by resolution. The compounds may, for example, be resolved into their components enantiomers by standard techniques, such as the formation of diastereomeric pairs by salt formation with an optically active acid, such as (-)-di-p-toluoyl-d-tartaric acid and/or (+)-di-p-toluoyl-l-tartaric acid followed by fractional crystallization and 25 regeneration of the free base. The compounds may also be resolved by formation of diastereomeric esters or amides, followed by chromatographic separation and removal of the chiral auxiliary. Alternatively, the compounds may be resolved using a chiral HPLC column.

30

During any of the processes for preparation of the compounds of the present invention, it may be necessary and/or desirable to protect sensitive or

ORT-1238

- reactive groups on any of the molecules concerned. This may be achieved by means of conventional protecting groups, such as those described in Protective Groups in Organic Chemistry, ed. J.F.W. McOmie, Plenum Press, 1973; and T.W. Greene & P.G.M. Wuts, Protective Groups in Organic
- 5 Synthesis, John Wiley & Sons, 1991. The protecting groups may be removed at a convenient subsequent stage using methods known from the art.

The method of treating PAR-1 mediated disorders (e.g., thrombotic disorders) described in the present invention may also be carried out using a

10 pharmaceutical composition comprising any of the compounds as defined herein and a pharmaceutically acceptable carrier. The pharmaceutical composition may contain between about 0.01 mg to about 100 mg, preferably from about 5 to about 50 mg, of the compound, and may be constituted into any form suitable for the mode of administration selected. Carriers include

15 necessary and inert pharmaceutical excipients, including, but not limited to, binders, suspending agents, lubricants, flavorants, sweeteners, preservatives, dyes, and coatings. Compositions suitable for oral administration include solid forms, such as pills, tablets, caplets, capsules (each including immediate release, timed release and sustained release formulations), granules, and

20 powders, and liquid forms, such as solutions, syrups, elixers, emulsions, and suspensions. Forms useful for parenteral administration include sterile solutions, emulsions and suspensions.

Advantageously, compounds of the present invention may be

25 administered in a single daily dose, or the total daily dosage may be administered in divided doses of two, three or four times daily. Furthermore, compounds for the present invention can be administered in intranasal form via topical use of suitable intranasal vehicles, or via transdermal skin patches well known to those of ordinary skill in that art. To be administered in the form of a

30 transdermal delivery system, the dosage administration will, of course, be continuous rather than intermittent throughout the dosage regimen.

ORT-1238

For instance, for oral administration in the form of a tablet or capsule, the active drug component can be combined with an oral, non-toxic pharmaceutically acceptable inert carrier such as ethanol, glycerol, water and the like. Moreover, when desired or necessary, suitable binders; lubricants, 5 disintegrating agents and coloring agents can also be incorporated into the mixture. Suitable binders include, without limitation, starch, gelatin, natural sugars such as glucose or beta-lactose, corn sweeteners, natural and synthetic gums such as acacia, tragacanth or sodium oleate, sodium stearate, magnesium stearate, sodium benzoate, sodium acetate, sodium chloride and 10 the like. Disintegrators include, without limitation, starch, methyl cellulose, agar, bentonite, xanthan gum and the like.

The liquid forms in suitably flavored suspending or dispersing agents such as the synthetic and natural gums, for example, tragacanth, acacia, 15 methyl cellulose and the like. For parenteral administration, sterile suspensions and solutions are desired. Isotonic preparations which generally contain suitable preservatives are employed when intravenous administration is desired.

20 The compound of the present invention can also be administered in the form of liposome delivery systems, such as small unilamellar vesicles, large unilamellar vesicles, and multilamellar vesicles. Liposomes can be formed from a variety of phospholipids, such as cholesterol, stearylamine or phosphatidylcholines.

25 Compounds of the present invention may also be delivered by the use of monoclonal antibodies as individual carriers to which the compound molecules are coupled. The compounds of the present invention may also be coupled with soluble polymers as targetable drug carriers. Such polymers can include 30 polyvinylpyrrolidone, pyran copolymer, polyhydroxypropylmethacrylamide-phenol, polyhydroxyethylaspartamidephenol, or polyethyleneoxidepolylysine substituted with palmitoyl residue. Furthermore, the compounds of the present

ORT-1238

invention may be coupled to a class of biodegradable polymers useful in achieving controlled release of a drug, for example, polylactic acid, polyepsilon caprolactone, polyhydroxy butyric acid, polyorthoesters, polyacetals, polydihydropyrans, polycyanoacrylates and cross-linked or amphipathic block copolymers of hydrogels.

Compounds of this invention may be administered in any of the foregoing compositions and according to dosage regimens established in the art whenever treatment of PAR-1 mediated disorders is required.

The daily dosage of the products may be varied over a wide range from about 0.01 mg to about 1,000 mg per adult human per day. For oral administration, the compositions are preferably provided in the form of tablets containing about 0.01, 0.05, 0.1, 0.5, 1.0, 2.5, 5.0, 10.0, 15.0, 25.0, 50.0, 100, 150, 200, 250 and 500 mg of the active ingredient for the symptomatic adjustment of the dosage to the patient to be treated. An effective amount of the drug is ordinarily supplied at a dosage level of from about 0.01 mg/kg to about 100 mg/kg of body weight per day. Preferably, the range is from about 0.03 mg/kg to about 10 mg/kg of body weight per day. The compounds may be administered on a regimen of about 1 time to about 4 times per day.

Optimal dosages to be administered may be readily determined by those skilled in the art, and will vary with the particular compound used, the mode of administration, the strength of the preparation, the mode of administration, and the advancement of the disease condition. In addition, factors associated with the particular patient being treated, including patient age, weight, diet and time of administration, will result in the need to adjust dosages.

Biology

The compounds of the present invention are thrombin receptor (PAR-1) antagonists. The compounds interrupt platelet activation induced by

ORT-1238

thrombin's proteolytic cleavage of its platelet surface receptor, and thereby inhibit platelet aggregation. Such compounds are, therefore, useful in treating platelet-mediated thrombotic disorders (e.g., arterial and venous thrombosis, acute myocardial infarction, reocclusion following thrombolytic therapy and angioplasty, and a variety of vaso-occlusive disorders) and other PAR-1 mediated disorders.

In Vitro Thrombin Receptor Binding Assay

CHRF membranes (Jones, *Biochim. Biophys. Acta* **1992**, 1136, 272) are thawed from -70°C, centrifuged at maximum speed for 5 min, washed twice with binding buffer (50 mM HEPES containing 5 mM MgCl₂ and 0.1% BSA), and re-suspended in binding buffer (25 µg/100 mL). 100 µL of membranes are added to the 24-Wallac plates and delivered to the Tomtech apparatus. In a typical experiment, 6 µL of samples (from a 125 µg/mL intermediary plate, 20% DMSO) and 44 µL buffer are delivered to the plates (final conc. of compounds is 3.7 µg/mL, 0.6% DMSO). Similarly, 6 µL 20% DMSO and 44 µL buffer are delivered to both column 1 (NSB) and column 12 (TB). 10 µL Ser-pFPhe-Har-Leu-Har-Lys-Tyr-NH₂ (721-40; 500 µM in deionized water) is added to column 1. 50 µL tritiated 721-40 (specific activity 46 Ci/mmol) is added to all the wells. The plates are mixed well for 20 seconds, incubated for 30 min, and then harvested with 10 mM HEPES/138 mM NaCl using the Skatron harvester. The filters (GF/C Brandel FPXLR 296) are presoaked 3 h in 0.5% polyethylenimine in HEPES/0.1M N-acetylglucosamine) are set in saran wrap and dried for 3 min in the microwave, and placed in sample bags (Wallac 145C-432). 4.5 mL scintillation fluid (Wallac, Betaplate Scint 1205-440) is added. The bags are sealed, placed in filter cassettes (Wallac 1450-104), and analyzed on the microbeta counter.

In Vitro Inhibition Of Thrombin-Induced Gel-Filtered Platelet Aggregation Assay

The percentage of platelet aggregation is calculated as an increase in light transmission of compound-treated platelet concentrate vs. control-treated platelet concentrate. Human blood is obtained from drug free, normal donors

ORT-1238

in tubes containing 0.13M sodium citrate. Platelet rich plasma (PRP) is collected by centrifugation of whole blood at 200 x g for 10 min at 25°C. The PRP (5 mL) is gel filtered through Sepharose 2B (bed volume 50 mL), and the platelet count is adjusted to 2×10^7 platelets per sample. The following constituents are added to a siliconized cuvette: concentrated platelet filtrate and Tyrode's buffer (0.14M NaCl, 0.0027M KCl, 0.012M NaHCO₃, 0.76 mM Na₂HPO₄, 0.0055M glucose, 2 mg/mL BSA and 5.0 mM HEPES @ pH 7.4) in an amount equal to 350 μ L, 50 μ L of 20 mM calcium and 50 μ L of the test compound. Aggregation is monitored in a BIODATA aggregometer for the 3 min following the addition of agonist (thrombin 50 μ L of 1 unit/mL).

Table 1 shows the biological activity of the compounds of the present invention. Table 1 contains IC₅₀ values (μ M) of the compounds against platelet aggregation stimulated by thrombin and IC₅₀ values (μ M) in a thrombin receptor (PAR-1) binding assay.

Examples

General Procedures: Resins and protected amino acids were purchased from Novabiochem, Bachem Bioscience, Advanced ChemTech or Synthe Tech. All other chemicals were obtained from commercial suppliers and used without further purification. ¹H and ¹³C NMR spectra were recorded on a Bruker AC 300B (300 MHz proton) or a Bruker AM-400 (400 MHz proton) spectrometer with Me₄Si as an internal standard (s = singlet, d = doublet, t = triplet, br = broad). APCI-MS and ES-MS were recorded on a VG Platform II mass spectrometer; methane was used for chemical ionization, unless noted otherwise. Accurate mass measurements were obtained by using a VG ZAB 2-SE spectrometer in the FAB mode. TLC was performed with Whatman 250- μ m silica gel plates. Preparative TLC was performed with Analtech 1000- μ m silica gel GF plates. Flash column chromatography was conducted with flash column silica gel (40-63 μ m) and column chromatography was conducted with standard silica gel. HPLC separations were carried out on three Waters PrepPak® Cartridges (25 x 100 mm, Bondapak® C18, 15-20 μ m, 125 Å) connected in series; detection was at 254 nm on a Waters 486 UV detector. Analytical HPLC was carried out on a Supelcosil ABZ+PLUS column (5 cm x 2.1 mm), with detection at 254 nm on a Hewlett Packard 1100 UV detector. Microanalysis was performed by Robertson Microlit Laboratories, Inc.

ORT-1238

In the examples and throughout this application, the following abbreviations have the meanings recited hereinafter:

| | | |
|----|-------|---|
| 5 | Ac | Acetyl |
| | ACN | Acetonitrile |
| | Bn | Benzyl |
| | Boc | t-Butoxycarbonyl |
| | DCC | 1,3-Dicyclohexylcarbodiimide |
| 10 | DCM | Dichloromethane |
| | DIC | Diisopropylcarbodiimide |
| | DIEA | Diisopropylethylamine |
| | DMF | N, N-Dimethylformamide |
| | Et | Ethyl |
| 15 | EtOAc | Ethyl acetate |
| | Fmoc | 9-Fluorenylmethoxycarbonyl |
| | h | Hour |
| | HBTU | 2-(1H-Benzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate |
| 20 | HOAc | Acetic acid |
| | HOBT | Hydroxybenzotriazole |
| | Me | Methyl |
| | min | Minute |
| | rt | room temperature |
| 25 | THF | Tetrahydrofuran |
| | TFA | Trifluoroacetic acid |
| | TLC | Thin layer chromatography |

Example 1

30 Synthesis of Compound 1 (Scheme AA)

Benzenepropanamide, N-[(1S)-3-amino-1-[(phenylmethyl)amino]
carbonyl]propyl]- α -[[[1-(2,6-dichlorophenyl)methyl]-3-
(1-pyrrolidinylmethyl)-1H-indazol-6-yl]amino]carbonyl]amino]-
35 3,4-difluoro-, (α S)- (Compound 1)

ORT-1238

6-Nitroindole (**AA1**, 1.0 g, 6.2 mmol) was suspended in a solution of sodium nitrite (4.3 g, 62 mmol) in H₂O (123 mL). To the suspension 6N HCl was added slowly until the pH was about pH 1. The resulting mixture was stirred at about rt, with protection from light, for about 2.5 h and extracted with EtOAc (120 mL X 3). The combined extracts were washed with H₂O (50 mL), brine (50 mL), dried (Na₂SO₄) and evaporated to give 1.13 g of indazole **AA2** as a yellow-pink solid. A solution of **AA2** (450 mg, 2.4 mmol) and pyrrolidine (836 mg, 11.8 mmol) in ClCH₂CH₂Cl:DMF:HOAc (90:9:1) were stirred at about rt for about 20 min, to which was then added NaB(OAc)₃H (1.25 g, 5.9 mmol) in one portion. The mixture was stirred at about rt for about 1 h, and then diluted with EtOAc (200 mL), washed with saturated NaHCO₃ (30 mL), brine (30 mL), dried (Na₂SO₄) and evaporated to afford 570 mg of **AA3** as a viscous brown solid. 6-Nitroindazole **AA3** (4.4 g, 17.8 mmol) was dissolved in dry THF (200 mL) under argon and 2,6-dichlorobenzyl bromide (4.3 g, 17.8 mmol) was added, followed by portionwise addition of pulverized KOH (1.17 g, 17.8 mmol) over about the next 20 min. The reaction was stirred at about rt for about 1 h and then evaporated *in vacuo* to an oil, which was partitioned between ethyl acetate (500 mL) and water (100 mL). The organic layer was separated and washed twice with water, three times with brine, dried (Na₂SO₄) and evaporated *in vacuo* to a brown solid. This was purified by flash column chromatography using DCM:MeOH (19:1) to afford 2.4 g of a tan solid which was combined with ferric chloride hexahydrate (0.30 g, 1.1 mmol) and activated charcoal (3.0 g, 0.25 mmol) in MeOH (200 mL). Dimethyl hydrazine (32 g, 0.53 mmol) was added and the reaction was refluxed for about 2 h, cooled to about rt and filtered through dicalite, which was washed several times with DCM:MeOH (4:1). The combined filtrates were evaporated *in vacuo* to a brown solid, which was purified by flash chromatography with DCM:MeOH:NH₄OH (90:8:1) to give 1.75 g of amine **AA4**.

Fmoc- α -N-Boc- γ -N-diaminobutyric acid **AA5** (10.8 g, 24.5 mmol) was stirred in ACN (300 mL) under argon as HOBt (3.75 g, 24.4 mmol) was added, followed by benzylamine (2.6 g, 24.3 mmol). DCC (10.4 g, 48.7 mmol) was added and the reaction was stirred at about rt for about 3 h, whereupon the resulting white solid was filtered and washed with cold ACN (14.4 g). The solid was stirred in ACN (500 mL) containing diethyl amine (25 mL) for about 2 h and a little solid was filtered; the filtrate was evaporated *in vacuo* to an oil, which was triturated three times with hexane (400 mL each) to a white solid.

ORT-1238

The solid was dissolved in ACN (400 mL) and HOBT (2.9 g, 19.1 mmol) and Fmoc-3,4 difluorophenylalanine (8.1 g, 19.1 mmol) were added, followed by DIC (4.81 g, 38.2 mmol) and stirred at about rt for about 16 h. The reaction was cooled in an ice bath and the white solid was filtered and washed with cold ACN. The solid was stirred in ACN (350 mL) containing diethylamine (35 mL) for about 5 h and evaporated *in vacuo* to a white solid, which was triturated three times with hexane, dissolved in chloroform (250 mL), dried (Na₂SO₄) and evaporated *in vacuo* to a white solid **AA6** (8.0 g).

6-Aminoindazole **AA4** (1.9 g, 5.0 mmol) and diisopropyl ethylamine (3.2 g, 25 mmol) in DCM (225 mL) under argon were cooled to about -20°C with CCl₄ / dry ice bath; 4-nitrophenyl chloroformate (1.10 g, 5.5 mmol) dissolved in DCM (10 mL) was added and the reaction was stirred at about -20°C for about 30 min. The dipeptide **AA6** (3.05 g, 5.0 mmol) was added and after about 30 min the reaction was allowed to warm to about rt and stirred for about an additional 6 h. The solution was ice bath cooled; a yellow solid was filtered and then washed with fresh, cold DCM. The solid was added to a solution (100 mL) of DCM:TFA:anisole (50:50:1), stirred at about rt for about 2.0 h and then evaporated *in vacuo* to a solid, which was triturated with diethyl ether (4X). The solid was purified by flash column chromatography using DCM:MeOH:NH₄OH (80:16:2) to give compound **1** as a white solid. The product was converted to the hydrochloride salt by dissolution in ACN and 1N HCl (20 mL, 1:1); evaporation *in vacuo* (3 X) and then lyophilization overnight afforded the white flaky solid Compound **1**•HCl: ¹H NMR (CD₃OD) δ 7.99 (s, 1 H), 7.72 (d, *J* = 8.4 Hz, 1 H), 7.45 - 7.05 (m, 12 H), 5.62 (s, 2 H), 4.64 (s, 2 H), 4.50 (m, 2 H), 4.39 (d, *J* = 2.6 Hz, 2 H), 3.52 (m, 2 H), 3.30 - 2.95 (m, 6 H), 2.30 - 1.85 (m, 6 H); ES-MS *m/z* 791 (MH⁺); Anal. Calc. C₄₀H₄₂Cl₂F₂N₈O₃•2HCl•2H₂O (900.68): C, 53.34; H, 5.37; N, 12.44; Cl, 15.74; KF, 4.00. Found: C, 53.15; H, 5.45; N, 12.38; Cl, 15.89; KF, 3.62.

Example 2

Synthesis of Compound 2 (Scheme AB)

5 Benzenepropanamide, N-[(1S)-3-amino-1-[(phenylmethyl)amino]
carbonyl]propyl)- α -[[[1-(2,6-dichlorophenyl)methyl]-3-
(1-pyrrolidinylmethyl)-1H-indazol-6-yl]amino]carbonyl]amino]-
4-chloro-, (α S)- (Compound 2)

To a solution of N- α -Fmoc-N- γ -Boc-diaminobutyric acid (**AB1**, 4.0 g, 9.1 mmol), BnNH₂ (1.07 g, 10 mmol) in CH₃CN (150 mL) was added HOBt (1.85 g, 13.7 mmol) and DCC (2.82 g, 13.7 mmol). The mixture was stirred at rt for 2.5 h, at which time TLC indicated that reaction was complete. The resulting white precipitates were collected by filtering and washing with CH₃CN to give 5.0 g of product (a mixture of the desired product and dicyclohexylurea). The combined filtrates were concentrated under vacuo and the residue was dissolved in EtOAc (150 mL). The solution was washed with saturated NaHCO₃, H₂O, brine, dried (Na₂SO₄), and evaporated to give a white powder which was recrystallized from CH₃CN to afford an additional product (1.7 g). The combined crude products were treated with 50% TFA in CH₂Cl₂ (80 mL) at rt for 1 h. The volatiles were removed under vacuo, and the residue was triturated with Et₂O to give **AB2** as a colorless solid (6.3 g). ¹H NMR showed it was a mixture **AB2** and dicyclohexylurea (ration 1:1.4). To a solution of the crude **AB2** (6.16 g, 7.14 mmol) and DIEA (2.71 g, 21.0 mmol) in DCM-DMF (1:1, 120 mL) was added 2-chlorotrityl chloride resin (4.0 g, 4.2 mmol) and the suspension was stirred at ambient temperature for 20 h. The reaction mixture was filtered on a sintered glass funnel and washed with DMF (2X), MeOH (3X), DCM (3X) and dried *in vacuo* to give resin (5.0 g). 4.9 g of resin was treated with 20% piperidine in DMF (80 mL) at rt for 2 h and then filtered, washed with DMF (2X), MeOH (2X), DCM (2X), Et₂O (2X) and dried *in vacuo* to afford resin **AB3** (4.15 g, loading level = 0.81 mmol/g, based on the mass loss during removing Fmoc group). A portion of **AB3** (1.1 g, 0.89 mmol) was suspended in DMF (30 mL) and treated with Fmoc-4-Cl-Phe-OH (0.94 g, 2.2 mmol), HOBt (0.30 g, 2.2 mmol), DIEA (0.58 g, 4.5 mmol), and HBTU (0.85 g, 2.2 mmol). The suspension was stirred at rt for 20 h and then filtered, washed with DMF, MeOH and DCM. The resulting resin was treated with 20% piperidine in DMF (30 mL) at rt for 2 h and then filtered, washed with DMF (2X), MeOH (2X), DCM (2X) and Et₂O (2X) to afford resin **AB4** (1.24 g). 6-Aminoindazole **AA4**

ORT-1238

(30 mg, 0.08 mmol) and diisopropyl ethylamine (52 mg, 0.4 mmol) in DCM (3 mL) under argon were cooled to about -20°C with CCl₄ / dry ice bath; 4-nitrophenyl chloroformate (17 mg, 0.085 mmol) dissolved in DCM (1 mL) was added dropwise and the reaction was stirred at about -20°C for about 15 min.

- 5 The resin-bound dipeptide amine **AB4** (47 mg, 0.034 mmol) was added and after about 20 min the reaction was allowed to warm to about rt and stirred for about an additional 2.5 h. The suspension was filtered and washed with MeOH, DCM and Et₂O and dried *in vacuo* to give resin **AB5**, which was then cleaved with TFA/DCM/anisole (50:50:1, 5 mL) at rt for 1.5 h, and reaction
10 mixture was filtered and washed with fresh 30% TFA in DCM. The filtrates were combined and evaporated *in vacuo*, and the residue was purified by preparative TLC using DCM-MeOH-28% NH₄OH (80:17:3) to give **2** as a gray solid: ¹H NMR (CD₃OD) δ 7.91 (s, 1 H), 7.68 (d, *J* = 8.8 Hz, 1 H), 7.45 - 7.17 (m, 12 H), 7.00 (d, *J* = 7.6 Hz, 1 H), 5.59 (s, 2 H), 4.55 - 4.36 (m, 6 H), 3.17 -
15 2.96 (m, 8 H), 2.21 - 1.90 (m, 6 H); ES-MS *m/z* 789 (MH⁺).

Example 3

Synthesis of Compound 3 (Scheme AC)

- 20 L-Alaninamide, 3,4-difluoro-*N*-[[[1-[(2,6-dichlorophenyl)methyl]-3-(1-pyrrolidinylmethyl)-1*H*-indazol-6-yl]amino]carbonyl]-L-phenylalanyl-*N*-(2-aminoethyl)-3-(4-pyridinyl)- (Compound 3)

- 2-Chlorotrityl chloride resin (4.8 g, 8.65 mmol; Advanced ChemTech) was
25 stirred in DMF (100 mL) as ethylene diamine **AC1** (15.6 g, 260 mmol) was added in and reaction stirred at ambient temperature for 16 h. The resin **AC2** was filtered on a sintered glass funnel and washed with DMF (4X), MeOH (3X), and DCM (3X) and dried *in vacuo*. A portion of resin **AC2** (2.0 g, 3.5 mmol) was placed in a solid phase hour-glass reactor and agitated (nitrogen bubbling)
30 in DMF (40 mL) with Fmoc-4-pyridyl alanine (3.9 g, 10 mmol), HOBT (1.53 g, 10 mmol), and DIC (1.26 g, 10 mmol) for 16 h. The solution was drawn off and the resin was washed with DMF (4X), DCM (4X) and DMF (2X) and then combined with 20% piperidine in DMF (25 mL) and agitated for 1.5 h. The solution was drained and the resin **AC3** was washed with DMF (5X) and
35 agitated in DMF (20mL) with Fmoc-3,4-diF-Phe-OH (4.23 g, 10 mmol), HOBT (1.53 g, 10 mmol) and DIC (1.26 g, 10 mmol) at ambient temperature for 16 h. The solution was removed and the resin was washed with DMF (5X), MeOH

ORT-1238

(3X), DCM (3X) and DMF (2X) and then combined with 20% piperidine in DMF (25 mL) and agitated for 1 h. The solution was drained and the resin was washed with DMF (4X), and DCM (4X) and dry DCM (3X) and stored *in vacuo* to give **AC4**. 6-Aminoindazole **AA4** (75 mg, 0.20 mmol) and diisopropyl ethylamine (145 mg, 1.1 mmol) in DCM (7 mL) under argon were cooled to about -20°C with CCl₄ / dry ice bath; 4-nitrophenyl chloroformate (38 mg, 0.19 mmol) dissolved in DCM (2 mL) was added dropwise and the reaction was stirred at about -20°C for about 20 min. The resin-bound dipeptide amine **AC4** (110 mg) was added and after about 20 min the reaction was allowed to warm to about rt and stirred for about an additional 18 h. The suspension was filtered and washed with MeOH, DCM and Et₂O and dried *in vacuo* to give resin **AC5**, which was then cleaved with TFA/DCM/anisole (50:50:1, 6 mL) at rt for 1 h, and reaction mixture was filtered and washed with fresh 30% TFA in DCM. The filtrates were combined and evaporated *in vacuo*, and the residue was purified by preparative TLC using DCM-MeOH-28% NH₄OH (85:12:3) to give compound **3** as a gray solid: ES-MS *m/z* 792 (MH⁺).


Example 4

As a specific embodiment of an oral composition, 100 mg of the Compound **1** of Example 1 is formulated with sufficient finely divided lactose to provide a total amount of about 580 mg to about 590 mg to fill a size O hard gel capsule.

While the foregoing specification teaches the principles of the present invention, with examples provided for the purpose of illustration, it will be understood that the practice of the invention encompasses all of the usual variations, adaptations and/or modifications as come within the scope of the following claims and their equivalents.

WHAT IS CLAIMED IS:

1. A compound of the following formula (I):



(I)

wherein:

A_1 and A_2 are each independently a D- or L-amino acid selected from the group consisting of alanine, β -alanine, arginine, homoarginine, cyclohexylalanine, citrulline, cysteine (optionally substituted with C_1 - C_4 alkyl, aryl, or arC_1 - C_4 alkyl), 2,4-diaminobutyric acid (optionally substituted with acyl, C_1 - C_4 alkyl, aroyl, amidino, or $MeC(NH)-$), 2,3-diaminopropionic acid (optionally substituted with acyl, C_1 - C_4 alkyl, aroyl, amidino, or $MeC(NH)-$), glutamine, glycine, indanylglycine, lysine (optionally substituted with acyl, C_1 - C_4 alkyl, aroyl, $MeC(NH)-$), valine, methionine, proline, serine (optionally substituted with C_1 - C_4 alkyl, aryl, or arC_1 - C_4 alkyl), homoserine (optionally substituted with C_1 - C_4 alkyl, aryl, or arC_1 - C_4 alkyl), tetrahydroisoquinoline-3-COOH, threonine (optionally substituted with C_1 - C_4 alkyl, aryl, or arC_1 - C_4 alkyl), ornithine (optionally substituted with acyl, C_1 - C_4 alkyl, aroyl, $MeC(NH)-$), and an unsubstituted or substituted aromatic amino acid selected from the group consisting of phenylalanine, heteroarylalanine, naphthylalanine, homophenylalanine, histidine, tryptophan, tyrosine, arylglycine, heteroarylglycine, aryl- β -alanine, and heteroaryl- β -alanine wherein the substituents on the aromatic amino acid are independently selected from one or more of halogen, C_1 - C_4 alkyl, C_1 - C_4 alkoxy, hydroxy, C_1 - C_4 alkoxycarbonyl,

ORT-1238

amino, amidino, guanidino, fluorinated C₁-C₄ alkyl, fluorinated C₁-C₄ alkoxy, C₁-C₄ alkylsulfonyl, C₁-C₄ alkylcarbonyl, cyano, aryl, heteroaryl, arC₁-C₄ alkyl, C₂-C₄ alkenyl, alkynyl, or nitro;

- 5 R₁ is selected from amino, C₁-C₈ alkylamino, C₁-C₈ dialkylamino, arylamino, arC₁-C₈ alkylamino, C₃-C₈ cycloalkylamino, heteroalkylC₁-C₈ alkylamino, heteroalkylC₁-C₈ alkyl-N-methylamino, C₁-C₈ dialkylaminoC₁-C₈ alkylamino, -N(C₁-C₈alkyl)-C₁-C₈ alkyl-N(C₁-C₈alkyl)₂, N(C₁-C₈alkyl)(C₁-C₈alkenyl), -N(C₁-C₈alkyl)(C₃-C₈cycloalkyl), heteroalkyl or substituted heteroalkyl wherein the substituent on the heteroalkyl is selected from oxo, amino, C₁-C₈ alkoxyC₁-C₈ alkyl, C₁-C₈ alkylamino or C₁-C₈ dialkylamino;

- 15 R₂ and R₃ are each independently selected from hydrogen, C₁-C₈ alkyl, C₃-C₈ cycloalkyl, C₃-C₈ cycloalkylC₁-C₈ alkyl, aryl, heteroalkyl, substituted heteroalkyl (wherein the substituent on the heteroalkyl is one or more substituents independently selected from C₁-C₈ alkoxy carbonyl, C₁-C₈ alkyl, or C₁-C₄ alkylcarbonyl), heteroalkylC₁-C₈ alkyl, indanyl, acetamidinoC₁-C₈ alkyl, aminoC₁-C₈ alkyl, C₁-C₈ alkylaminoC₁-C₈ alkyl, C₁-C₈ dialkylaminoC₁-C₈ alkyl, 20 unsubstituted or substituted heteroarylC₁-C₈ alkyl or unsubstituted or substituted arC₁-C₈ alkyl, wherein the substituent on the aralkyl or heteroarylalkyl group is one or more substituents independently selected from halogen, nitro, amino, C₁-C₈ alkyl, C₁-C₈ alkoxy, hydroxy, cyano, C₁-C₄ alkylcarbonyl, C₁-C₈ alkoxy carbonyl, hydroxyC₁-C₈ alkyl or aminosulfonyl; or

- 25 R₂ and R₃, together with the nitrogen to which they are attached, alternatively form an unsubstituted or substituted heteroalkyl group selected from piperidinyl, piperazinyl, morpholinyl or pyrrolidinyl, wherein the substituent is one or more substituents independently selected from C₁-C₈ alkyl C₁-C₈ alkoxy carbonyl or C₁-C₄ alkylcarbonyl;

- 35 R₄ is selected from unsubstituted or substituted aryl, arC₁-C₈ alkyl, C₃-C₈ cycloalkyl, or heteroaryl, where the substituents on the aryl, arC₁-C₈ alkyl, cycloalkyl or heteroaryl group are independently selected from one or more of halogen, nitro, amino, cyano, hydroxyalkyl, C₁-C₈ alkyl, C₁-C₈ alkoxy, hydroxy,

ORT-1238

C₁-C₄ alkylcarbonyl, C₁-C₈ alkoxy carbonyl, fluorinated C₁-C₄ alkyl, fluorinated C₁-C₄ alkoxy, C₁-C₄ alkylsulfonyl;

R₅ is selected from hydrogen or C₁-C₈ alkyl;

X is oxygen or sulfur;

m is an integer selected from 0, 1, 2 or 3;

n is an integer selected from 1 or 2; and

p is an integer selected from 0 or 1;

and pharmaceutically acceptable salts thereof.

2. The compound of Claim 1, wherein:

A₁ and A₂ are each independently an L-amino acid selected from the group consisting of alanine, β-alanine, arginine, homoarginine, cyclohexylalanine, citrulline, cysteine (optionally substituted with C₁-C₄ alkyl, aryl, or arC₁-C₄ alkyl), 2,4-diaminobutyric acid (optionally substituted with acyl, C₁-C₄ alkyl, aroyl, amidino, or MeC(NH)-), 2,3-diaminopropionic acid (optionally substituted with acyl, C₁-C₄ alkyl, aroyl, amidino, or MeC(NH)-), glutamine, glycine, indanylglycine, lysine (optionally substituted with acyl, C₁-C₄ alkyl, aroyl, MeC(NH)-), valine, methionine, proline, serine (optionally substituted with C₁-C₄ alkyl, aryl, or arC₁-C₄ alkyl), homoserine (optionally substituted with C₁-C₄ alkyl, aryl, or arC₁-C₄ alkyl), tetrahydroisoquinoline-3-COOH, threonine (optionally substituted with C₁-C₄ alkyl, aryl, or arC₁-C₄ alkyl), ornithine (optionally substituted with acyl, C₁-C₄ alkyl, aroyl, MeC(NH)-), and an unsubstituted or substituted aromatic amino acid selected from the group consisting of phenylalanine, heteroarylalanine, naphthylalanine, homophenylalanine, histidine, tryptophan, tyrosine, arylglycine, heteroarylglycine, aryl-β-alanine, and heteroaryl-β-alanine wherein the substituents on the aromatic amino acid are independently selected from one or more of halogen, C₁-C₄ alkyl, C₁-C₄ alkoxy, hydroxy, C₁-C₄ alkoxy carbonyl, amino, amidino, guanidino, fluorinated C₁-C₄ alkyl, fluorinated C₁-C₄ alkoxy, C₁-

ORT-1238

C₄ alkylsulfonyl, C₁-C₄ alkylcarbonyl, cyano, aryl, heteroaryl, arC₁-C₄ alkyl, C₂-C₄ alkenyl, alkynyl, or nitro;

R₁ is selected from amino, C₁-C₆ alkylamino, C₁-C₆ dialkylamino, arylamino, arC₁-C₆ alkylamino, heteroalkylC₁-C₆ alkylamino, -N(C₁-C₆alkyl)-C₁-C₆ alkyl-N(C₁-C₆alkyl)₂, heteroalkyl or substituted heteroalkyl wherein the substituent on the heteroalkyl is selected from oxo, amino, C₁-C₆alkoxyC₁-C₆ alkyl, C₁-C₆ alkylamino or C₁-C₆ dialkylamino;

R₂ is selected from hydrogen or C₁-C₆ alkyl;

R₃ is selected from C₁-C₈ alkyl, C₃-C₆ cycloalkyl, C₃-C₆cycloalkylC₁-C₆alkyl, aryl, heteroarylC₁-C₆ alkyl, substituted heteroarylC₁-C₆alkyl wherein the substituent is C₁-C₄ alkyl, heteroalkyl, heteroalkylC₁-C₆ alkyl, indanyl, acetamidinoC₁-C₆ alkyl, aminoC₁-C₆ alkyl, C₁-C₆alkylaminoC₁-C₆ alkyl, C₁-C₆ dialkylaminoC₁-C₆ alkyl, arC₁-C₈alkyl, substituted arC₁-C₈ alkyl wherein the substituent on the aralkyl group is one to five substituents independently selected from halogen, nitro, amino, C₁-C₆ alkyl, C₁-C₆ alkoxy, C₁-C₆ alkoxycarbonyl, hydroxyalkyl or aminosulfonyl;

R₂ and R₃, together with the nitrogen to which they are attached, alternatively form an unsubstituted or substituted heteroalkyl group selected from piperidinyl, piperazinyl or pyrrolidinyl, wherein the substituent is independently one or two substituents selected from C₁-C₆ alkyl;

R₄ is selected from unsubstituted or substituted aryl, arC₁-C₆ alkyl, C₃-C₆cycloalkyl or heteroaryl, where the substituents on the aryl, aralkyl, cycloalkyl or heteroaryl group are independently selected from one to three substituents selected from halogen, cyano, C₁-C₄ alkyl, C₁-C₄ alkoxy, C₁-C₄alkoxycarbonyl, fluorinated C₁-C₄ alkyl, fluorinated C₁-C₄ alkoxy or C₁-C₄alkylsulfonyl;

R₅ is hydrogen;

X is oxygen; and

p is 1;

and pharmaceutically acceptable salts thereof.

3. The compound of Claim 2, wherein:

5

A₁ is an L-amino acid selected from the group consisting of alanine, arginine, cyclohexylalanine, glycine, proline, tetrahydroisoquinoline-3-COOH, and an unsubstituted or substituted aromatic amino acid selected from the group consisting of phenylalanine, naphthylalanine, homophenylalanine, and

10

O-methyl tyrosine, wherein the substituents on the aromatic amino acid are independently one to five substituents selected from halogen, C₁-C₄ alkyl, C₁-C₄ alkoxy, hydroxy, C₁-C₄ alkoxycarbonyl, amino, amidino, guanidino, fluorinated C₁-C₄ alkyl, fluorinated C₁-C₄ alkoxy, C₁-C₄ alkylsulfonyl, C₁-C₄ alkylcarbonyl, cyano, aryl, heteroaryl, arC₁-C₄ alkyl, C₂-C₄ alkenyl, alkynyl, or
15 nitro;

15

A₂ is an L-amino acid selected from the group consisting of alanine, β-alanine, arginine, citrulline, cysteine (optionally substituted with C₁-C₄ alkyl, aryl, or arC₁-C₄ alkyl), 2,4-diaminobutyric acid (optionally substituted with acyl, C₁-C₄ alkyl, aroyl, amidino, or MeC(NH)-), 2,3- diaminopropionic acid (optionally substituted with acyl, C₁-C₄ alkyl, aroyl, amidino, or MeC(NH)-), glutamine, glycine, lysine (optionally substituted with acyl, C₁-C₄ alkyl, aroyl, MeC(NH)-), valine, methionine, serine (optionally substituted with C₁-C₄ alkyl, aryl, or arC₁-C₄ alkyl), homoserine (optionally substituted with C₁-C₄ alkyl, aryl, or arC₁-C₄ alkyl), threonine (optionally substituted with C₁-C₄ alkyl, aryl, or arC₁-C₄ alkyl), ornithine (optionally substituted with acyl, C₁-C₄ alkyl, aroyl, MeC(NH)-), and an unsubstituted or substituted aromatic amino acid selected from the group consisting of phenylalanine, heteroarylalanine, and histidine, wherein the substituents of the aromatic amino acid are independently one to
20 five substituents selected from halogen, C₁-C₄ alkyl, C₁-C₄ alkoxy, hydroxy, C₁-C₄ alkoxycarbonyl, amino, amidino, guanidino, fluorinated C₁-C₄ alkyl, fluorinated C₁-C₄ alkoxy, C₁-C₄ alkylsulfonyl, C₁-C₄ alkylcarbonyl, cyano, aryl, heteroaryl, arC₁-C₄ alkyl, C₂-C₄ alkenyl, alkynyl, or nitro;

20

25

30

ORT-1238

R₂ is selected from hydrogen or C₁-C₄ alkyl; and

m and n are both 1;

5

and pharmaceutically acceptable salts thereof.

4. The compound of Claim 3, wherein:

10

A₁ is an L-amino acid selected from the group consisting of alanine, arginine, cyclohexylalanine, proline, tetrahydroisoquinoline-3-COOH, and an unsubstituted or substituted aromatic amino acid selected from the group consisting of phenylalanine, naphthylalanine, homophenylalanine, and O-methyl tyrosine, wherein the substituents on the aromatic amino acid are independently one to two substituents selected from halogen, C₁-C₄ alkyl, C₁-C₄ alkoxy, hydroxy, C₁-C₄ alkoxycarbonyl, amino, amidino, guanidino, fluorinated C₁-C₄ alkyl, fluorinated C₁-C₄ alkoxy, C₁-C₄ alkylsulfonyl, C₁-C₄ alkylcarbonyl, cyano, aryl, heteroaryl, arC₁-C₄ alkyl, C₂-C₄ alkenyl, alkynyl, or nitro;

15

20

A₂ is an L-amino acid selected from the group consisting of alanine, β-alanine, arginine, citrulline, cysteine (optionally substituted with C₁-C₄ alkyl, aryl, or arC₁-C₄ alkyl), 2,4-diaminobutyric acid (optionally substituted with acyl, C₁-C₄ alkyl, aroyl, amidino, or MeC(NH)-), 2,3-diaminopropionic acid (optionally substituted with acyl, C₁-C₄ alkyl, aroyl, amidino, or MeC(NH)-), glutamine, glycine, lysine (optionally substituted with acyl, C₁-C₄ alkyl, aroyl, MeC(NH)-), valine, methionine, serine (optionally substituted with C₁-C₄ alkyl, aryl, or arC₁-C₄ alkyl), homoserine (optionally substituted with C₁-C₄ alkyl, aryl, or arC₁-C₄ alkyl), threonine (optionally substituted with C₁-C₄ alkyl, aryl, or arC₁-C₄ alkyl), ornithine (optionally substituted with acyl, C₁-C₄ alkyl, aroyl, MeC(NH)-), and an unsubstituted or substituted aromatic amino acid selected from the group consisting of phenylalanine, heteroarylalanine, and histidine, wherein the substituents on the aromatic amino acid are independently one to two

25


30

ORT-1238

substituents selected from halogen, C₁-C₄ alkyl, C₁-C₄ alkoxy, hydroxy, C₁-C₄ alkoxycarbonyl, amino, amidino, guanidino, fluorinated C₁-C₄ alkyl, fluorinated C₁-C₄ alkoxy, C₁-C₄ alkylsulfonyl, C₁-C₄ alkylcarbonyl, cyano, aryl, heteroaryl, arC₁-C₄ alkyl, C₂-C₄ alkenyl, alkynyl, or nitro;

5

R₁ is selected from diethylamino, di-(*n*-propyl)amino,



10

R₂ is selected from hydrogen, methyl or ethyl;

R₃ is selected from 2-indanyl, phenyl, cyclohexylmethyl, cyclopentyl, pyridylmethyl, furanylmethyl, 2-(4-methyl-furanyl)methyl, thienylmethyl, diphenylmethyl, 4-imidazolylethyl, 2-(4-N-methyl)imidazolylethyl, *n*-octyl, phenyl-*n*-propyl, aminoethyl, aminopropyl, amino-*n*-pentyl, dimethylaminoethyl, 4-aminophenylsulfonylaminomethyl, acetamidineylethyl, 2-N-pyrrolidinylethyl, N-ethoxycarbonylpiperidinyl, unsubstituted or substituted phenylethyl and unsubstituted or substituted benzyl wherein the substituents on the phenylethyl or benzyl are independently one or two substituents selected from methyl, fluorine, chlorine, nitro, methoxy, methoxycarbonyl or hydroxymethyl; or

20

R₂ and R₃, together with the nitrogen to which they are attached, alternatively form a heteroalkyl group selected from piperidinyl, or 4-(N-methyl)piperazinyl; and

25


R₄ is selected from cyclohexyl, 2-naphthyl, phenylethyl, 4-fluorophenylethyl or unsubstituted or substituted phenyl, where the substituents on the phenyl are independently selected from one to two substituents selected from fluorine, chlorine, iodine, methyl, cyano or trifluoromethyl;

30

and pharmaceutically acceptable salts thereof.

5. The compound of Claim 4, wherein:


ORT-1238



and pharmaceutically acceptable salts thereof.

5

6. The compound of Claim 5, wherein:



and pharmaceutically acceptable salts thereof.

10

7. The compound of Claim 6, wherein:

A₁ is selected from 3,4-Difluorophenylalanine or 4-Chlorophenylalanine;

A₂ is selected from 2,4-Diaminobutyric acid or 4-Pyridylalanine;

R₂ is hydrogen; and

R₃ is selected from benzyl or 2-aminoethyl;

and pharmaceutically acceptable salts thereof.

20

8. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and a compound of Claim 1.

9. A pharmaceutical composition made by mixing a compound of Claim 1 and a pharmaceutically acceptable carrier.

25

10. A process for making a pharmaceutical composition comprising mixing a compound of Claim 1 and a pharmaceutically acceptable carrier.

ORT-1238

11. A method of treating a condition selected from the group consisting of thrombosis, restenosis, hypertension, heart failure, arrhythmia, myocardial infarction, glomerulonephritis, reocclusion following thrombolytic therapy, reocclusion following angioplasty, inflammation, angina, stroke, atherosclerosis, ischemic conditions, a vaso-occlusive disorder, neurodegenerative disorders, Angiogenesis related disorders and cancer in a subject in need thereof comprising administering to the subject a therapeutically effective amount of the compound of Claim 1.

12. The method of Claim 11, wherein the therapeutically effective amount of the compound is from about 0.1 mg/kg/day to about 300 mg/kg/day.

13. A method of treating a condition selected from the group consisting of thrombosis, restenosis, hypertension, heart failure, arrhythmia, myocardial infarction, glomerulonephritis, reocclusion following thrombolytic therapy, reocclusion following angioplasty, inflammation, angina, stroke, atherosclerosis, ischemic conditions, a vaso-occlusive disorder, neurodegenerative disorders, Angiogenesis related disorders and cancer in a subject in need thereof comprising administering to the subject a therapeutically effective amount of the composition of Claim 8.

14. The method of Claim 13, wherein the therapeutically effective amount of the compound is from about 0.1 mg/kg/day to about 300 mg/kg/day.

15. A method of inhibiting platelet aggregation in a subject in need thereof comprising administering to the subject a therapeutically effective amount of the compound of Claim 1.

16. The method of Claim 15, wherein the therapeutically effective amount of the compound is from about 0.1 mg/kg/day to about 300 mg/kg/day.

ORT-1238

17. A method of inhibiting platelet aggregation in a subject in need thereof comprising administering to the subject a therapeutically effective amount of the composition of Claim 8.

5 18. The method of Claim 17, wherein the therapeutically effective amount of the compound is from about 0.1 mg/kg/day to about 300 mg/kg/day.


19. A method of treating a condition mediated by thrombin receptor (PAR-1) in a subject in need thereof comprising administering to the subject a
10 therapeutically effective amount of the compound of Claim 1.

20. The method of Claim 19, wherein the therapeutically effective amount of the compound is from about 0.1 mg/kg/day to about 300 mg/kg/day.

15 21. A method of treating a condition mediated by thrombin receptor (PAR-1) in a subject in need thereof comprising administering to the subject a therapeutically effective amount of the composition of Claim 8.

22. The method of Claim 21, wherein the therapeutically effective amount of
20 the compound is from about 0.1 mg/kg/day to about 300 mg/kg/day.

23. A process for preparing a compound of the formula (II):



(II)

wherein:

ORT-1238

A_1 and A_2 are each independently a D- or L-amino acid selected from the group consisting of alanine, β -alanine, arginine, homoarginine, cyclohexylalanine, citrulline, cysteine (optionally substituted with C_1 - C_4 alkyl, aryl, or arC_1 - C_4 alkyl), 2,4-diaminobutyric acid (optionally substituted with acyl, C_1 - C_4 alkyl, aroyl, amidino, or $MeC(NH)-$), 2,3-diaminopropionic acid (optionally substituted with acyl, C_1 - C_4 alkyl, aroyl, amidino, or $MeC(NH)-$), glutamine, glycine, indanylglycine, lysine (optionally substituted with acyl, C_1 - C_4 alkyl, aroyl, $MeC(NH)-$), valine, methionine, proline, serine (optionally substituted with C_1 - C_4 alkyl, aryl, or arC_1 - C_4 alkyl), homoserine (optionally substituted with C_1 - C_4 alkyl, aryl, or arC_1 - C_4 alkyl), tetrahydroisoquinoline-3-COOH, threonine (optionally substituted with C_1 - C_4 alkyl, aryl, or arC_1 - C_4 alkyl), ornithine (optionally substituted with acyl, C_1 - C_4 alkyl, aroyl, $MeC(NH)-$), and an unsubstituted or substituted aromatic amino acid selected from the group consisting of phenylalanine, heteroarylalanine, naphthylalanine, homophenylalanine, histidine, tryptophan, tyrosine, arylglycine, heteroarylglycine, aryl- β -alanine, and heteroaryl- β -alanine wherein the substituents on the aromatic amino acid are independently selected from one or more of halogen, C_1 - C_4 alkyl, C_1 - C_4 alkoxy, hydroxy, C_1 - C_4 alkoxycarbonyl, amino, amidino, guanidino, fluorinated C_1 - C_4 alkyl, fluorinated C_1 - C_4 alkoxy, C_1 - C_4 alkylsulfonyl, C_1 - C_4 alkylcarbonyl, cyano, aryl, heteroaryl, arC_1 - C_4 alkyl, C_2 - C_4 alkenyl, alkynyl, or nitro;

R_1 is selected from amino, C_1 - C_8 alkylamino, C_1 - C_8 dialkylamino, arylamino, arC_1 - C_8 alkylamino, C_3 - C_8 cycloalkylamino, heteroalkyl C_1 - C_8 alkylamino, heteroalkyl C_1 - C_8 alkyl-N-methylamino, C_1 - C_8 dialkylamino C_1 - C_8 alkylamino, $-N(C_1-C_8alkyl)-C_1-C_8$ alkyl- $N(C_1-C_8alkyl)_2$, $N(C_1-C_8alkyl)(C_1-C_8alkenyl)$, $-N(C_1-C_8alkyl)(C_3-C_8cycloalkyl)$, heteroalkyl or substituted heteroalkyl wherein the substituent on the heteroalkyl is selected from oxo, amino, C_1 - C_8 alkoxy C_1 - C_8 alkyl, C_1 - C_8 alkylamino or C_1 - C_8 dialkylamino;

R_2 and R_3 are each independently selected from hydrogen, C_1 - C_8 alkyl, C_3 - C_8 cycloalkyl, C_3 - C_8 cycloalkyl C_1 - C_8 alkyl, aryl, heteroalkyl, substituted


ORT-1238

heteroalkyl (wherein the substituent on the heteroalkyl is one or more substituents independently selected from C₁-C₈ alkoxy, carbonyl, C₁-C₈ alkyl, or C₁-C₄ alkylcarbonyl), heteroalkylC₁-C₈ alkyl, indanyl, acetamidinoC₁-C₈ alkyl, aminoC₁-C₈ alkyl, C₁-C₈ alkylaminoC₁-C₈ alkyl, C₁-C₈ dialkylaminoC₁-C₈ alkyl, unsubstituted or substituted heteroarylC₁-C₈ alkyl or unsubstituted or substituted arC₁-C₈ alkyl, wherein the substituent on the aralkyl or heteroarylalkyl group is one or more substituents independently selected from halogen, nitro, amino, C₁-C₈ alkyl, C₁-C₈ alkoxy, hydroxy, cyano, C₁-C₄ alkylcarbonyl, C₁-C₈ alkoxy, carbonyl, hydroxyC₁-C₈ alkyl or aminosulfonyl; or


R₂ and R₃, together with the nitrogen to which they are attached, alternatively form an unsubstituted or substituted heteroalkyl group selected from piperidinyl, piperazinyl, morpholinyl or pyrrolidinyl, wherein the substituent is one or more substituents independently selected from C₁-C₈ alkyl, C₁-C₈ alkoxy, carbonyl, or C₁-C₄ alkylcarbonyl;

R₄ is selected from unsubstituted or substituted aryl, arC₁-C₈ alkyl, C₃-C₈ cycloalkyl, or heteroaryl, where the substituents on the aryl, arC₁-C₈ alkyl, cycloalkyl or heteroaryl group are independently selected from one or more of halogen, nitro, amino, cyano, hydroxyalkyl, C₁-C₈ alkyl, C₁-C₈ alkoxy, hydroxy, C₁-C₄ alkylcarbonyl, C₁-C₈ alkoxy, carbonyl, fluorinated C₁-C₄ alkyl, fluorinated C₁-C₄ alkoxy, C₁-C₄ alkylsulfonyl;

comprising reacting a compound of the formula **AAG6**:



with a compound of the formula **AAG4**:



in the presence of a phosgene equivalent to form the compound of formula (II).

ABSTRACT OF THE INVENTION

The invention is directed to novel indazole peptidomimetic compounds
5 which are useful as thrombin receptor antagonists for the treatment of diseases
associated with thrombosis, restenosis, hypertension, heart failure, arrhythmia,
inflammation, angina, stroke, atherosclerosis, ischemic conditions,
Angiogenesis related disorders, cancer, and neurodegenerative disorders.
Pharmaceutical compositions comprising the substituted indazole
10 peptidomimetics of the present invention and methods of treating conditions
mediated by the thrombin receptor are also disclosed.

DECLARATION AND POWER OF ATTORNEY FOR PATENT APPLICATION

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name,

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled NOVEL INDAZOLE PEPTIDOMIMETICS AS THROMBIN RECEPTOR ANTAGONISTS, the specification of which

(check one) ☒ is attached hereto.

☐ was filed on _____ as

Application Serial No. _____

and was amended on _____.
(if applicable)

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is material to patentability as defined in 37 CFR 1.56.

I hereby claim foreign priority benefits under Title 35, United States Code, §119 (a)-(d) or §365(b) of any foreign application(s) for patent or inventor's certificate, or §365(a) of any PCT international application which designated at least one country other than the United States of America, listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate, or of any PCT international application having a filing date before that of the application on which priority is claimed.

Prior Foreign Application(s):

| Country | Application Number | Date of Filing | Priority Claimed Under 35 U.S.C. 119 | |
|---------|--------------------|----------------|--------------------------------------|-----------------------------|
| | | | <input type="checkbox"/> YES | <input type="checkbox"/> NO |
| | | | <input type="checkbox"/> YES | <input type="checkbox"/> NO |
| | | | <input type="checkbox"/> YES | <input type="checkbox"/> NO |

I hereby claim the benefit under Title 35, United States Code, §119(e) of any United States provisional application(s) listed below:

60/141,553
(Application Number)

6/29/99
(Filing Date)

(Application Number)

(Filing Date)

I hereby claim the benefit under Title 35, United States Code, §120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, §112, I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, §1.56(a) which occurred between the filing date of the prior application and the national or PCT international filing date of this application:

Application Serial No. Filing Date Status

Application Serial No. Filing Date Status

I hereby appoint the following attorney(s) and/or agent(s) to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith as well as to file equivalent patent applications in countries foreign to the United States including the filing of international patent applications in accordance with the Patent Cooperation Treaty: Audley A. Ciamporzero, Jr. (Reg. #26,051), Steven P. Berman (Reg. #24,772), Andrea L. Colby (Reg. #30,194), Michael Stark (Reg. #32,495), and Hal B. Woodrow (Reg. #32,501) One Johnson & Johnson Plaza, New Brunswick, NJ 08933.

Address all telephone calls to Hal B. Woodrow at telephone no.
(732) 524-2976.

Address all correspondence to Audley A. Ciamporzero, Jr., One
Johnson & Johnson Plaza, New Brunswick, NJ 08933-7003.

I hereby declare that all statements made herein of my own
knowledge are true and that all statements made on information
and belief are believed to be true; and further that these
statements were made with the knowledge that willful false
statements and the like so made are punishable by fine or
imprisonment, or both, under Section 1001 of Title 18 of the
United States Code and that such willful false statements may
jeopardize the validity of the application or any patent issued
thereon.


Inventor's Signature:
Full Name of First Joint
Inventor


Han-Cheng Zhang

Date: 5/23/2000

Citizenship: P.R. of China
Residence: 109 Bryan Way, Lansdale, PA 19446
Post Office Address: Same as Above


Inventor's Signature:
Full Name of Second Joint
Inventor


Bruce E. Maryanoff

Date: 24 May 00

Citizenship: USA
Residence: 4029 Devonshire Drive, Forest Grove, PA 18922
Post Office Address: Same as Above


Inventor's Signature:
Full Name of Third Joint
Inventor


Anjali Pandey

Date: 6/6/2000

Citizenship: USA
Residence: 4506 Armians Avenue, Fremont, CA 94555
Post Office Address: Same as Above

Inventor's Signature:
Full Name of Fourth Joint
Inventor


Robert M. Scarborough

Date: 6/6/2000

Citizenship: USA

Residence: 22 Greenbrier Court, Half Moon Bay, CA 94019

Post Office Address: Same as Above